

The effects of early versus late time-restricted feeding on metabolic disease risk factors in adults at increased risk of developing type 2 diabetes: Is there an optimal time to eat?

Submission date 23/07/2019	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
Registration date 01/08/2019	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 12/12/2022	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

There is a rise in the number of people suffering from type 2 diabetes and a large proportion of the UK population are at increased risk of developing the condition. Some of the risk factors for type 2 diabetes include being overweight, having a family history of the condition, high blood pressure, abnormal blood cholesterol levels and ethnicity. This study seeks to investigate whether restricting dietary intake to a set number of hours in the day – either early in the day or late in the day – will make a difference to known risk factors for developing type 2 diabetes. By restricting dietary intake to a set number of hours in the day, this will increase the amount of time spent fasting (not eating) each day. This type of diet is known as time-restricted feeding. There is some evidence to suggest that time-restricted feeding may provide some health benefits in healthy individuals by helping to reduce the amount of energy consumed through food, lowering body weight and body fat, as well as improving markers of metabolic health. However, it is not yet known whether it is more beneficial to restrict our dietary intake to earlier or later in the day. Additionally, adherence to a change in diet is critical in predicting its success and long-term outcomes. However, when people find themselves in an environment with unlimited access to food during the day, it can be difficult to maintain successful behavioural changes. Therefore, the study will also look at how following this type of diet may impact on well-being, social life and the functioning of families (especially at mealtimes), to assess the suitability of the intervention for the general public, so that it does not negatively affect quality of life. Overall, this study will compare the effects of early versus late time-restricted feeding in adults at increased risk of developing type 2 diabetes.

Who can participate?

Adults aged 18-65 years old who are overweight

What does the study involve?

Participants are randomly allocated to one of three groups for 10 weeks:

1. A control group – where normal eating times are maintained

2. An early time-restricted feeding group – where eating is limited to between 7 am and 3 pm each day (\pm 1 hour)
 3. A late time-restricted feeding group – where eating is limited to between 12 pm and 8 pm each day (\pm 1 hour)
- Participants are asked to attend the University of Surrey for a total of five study day visits during this period for a variety of health assessments.

What are the possible benefits and risks of participating?

A variety of health assessments will be conducted as part of the study, including blood tests and scans to look at body fat distribution. All participants will receive one-to-one dietary support from a Registered Dietitian during the 10-week intervention period. There is a small risk of light bruising at the blood sample site, but all samples will be taken by trained phlebotomists the same way as during a blood test. Participants will also be asked to do an oral glucose tolerance test to look at how their body metabolises sugars. This will involve using a lancet device (used to collect blood samples) which will involve a short sharp prick to the tip of your finger. To measure the distribution of fat across the body, participants will receive a total of 2 dual-energy x-ray absorptiometry (DEXA) scans during the entire study. A DEXA scan exposes participants to ionising radiation, but the total radiation dose received from a DEXA is \sim 8 μ Sv per scan. This is equivalent to the amount of radiation that you would be exposed to if you spent 45 minutes on a transatlantic flight or the amount of radiation contained in 80 g of brazil nuts grown in Brazil. Although the dose of radiation participants are exposed to during the DEXA scan is minimal, there is a risk that even small doses of radiation can cause cancer.

Where is the study run from?
University of Surrey (UK)

When is the study starting and how long is it expected to run for?
May 2019 to June 2023

Who is funding the study?
University of Surrey (UK)

Who is the main contact?
1. Shantel Lynch
s.lynch@surrey.ac.uk
2. Dr Denise Robertson
m.robertson@surrey.ac.uk
3. Dr Jonathan Johnston
j.johnston@surrey.ac.uk

Contact information

Type(s)
Public

Contact name
Mrs Shantel Lynch

ORCID ID
<https://orcid.org/0000-0003-3117-7362>

Contact details

University of Surrey
Guildford
Surrey
United Kingdom
GU2 7XH
+44 (0)1483300800
s.lynch@surrey.ac.uk

Type(s)

Scientific

Contact name

Mrs Shantel Lynch

ORCID ID

<https://orcid.org/0000-0003-3117-7362>

Contact details

University of Surrey
Guildford
Surrey
United Kingdom
GU2 7XH
+44 (0)1483300800
s.lynch@surrey.ac.uk

Type(s)

Scientific

Contact name

Dr Denise Robertson

ORCID ID

<https://orcid.org/0000-0001-5461-9544>

Contact details

University of Surrey
Guildford
Surrey
United Kingdom
GU2 7XH
+44 (0)1483300800
m.robertson@surrey.ac.uk

Type(s)

Scientific

Contact name

Dr Jonathan Johnston

Contact details

University of Surrey
Guildford
Surrey
United Kingdom
GU2 7XH
+44 (0)1483300800
j.johnston@surrey.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

257999

Study information

Scientific Title

Early versus late time-restricted feeding: its effect on metabolic disease risk factors in adults at increased risk of developing type 2 diabetes

Study objectives

This study will test the primary hypothesis that 10 weeks of time-restricted feeding (TRF) will reduce insulin resistance and LDL-cholesterol in adults at increased risk of type 2 diabetes, as well as promote reductions in dietary energy intake, body weight and adiposity. The secondary hypothesis will be that early-TRF will be metabolically superior to late-TRF.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 26/04/2019, London – Surrey Research Ethics Committee (Whitefriars, Level 3, Block B, Lewins Mead, Bristol, BS1 2NT, UK; Tel: +44 (0)207 104 8310; Email: nrescommittee.secoast-surrey@nhs.net), REC ref: 19/LO/0540, IRAS: 257999

Study design

Three-arm parallel randomised controlled dietary intervention trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Type 2 diabetes mellitus

Interventions

After an initial 1-week baseline period, participants will be randomly assigned to one of three experimental groups for an intervention period of approximately 10 weeks.

The first (control) group will be asked to maintain their habitual sleep-wake and feed-fast routines throughout the intervention period.

The second, 'early-TRF', group will be asked to maintain their habitual sleep-wake routine, but restrict the duration of their eating times during the day to between 7am and 3pm (± 1 hour).

The third, 'late-TRF', group will also be asked to maintain their habitual sleep-wake routine, but restrict the duration of their eating times during the day to between 12pm and 8pm (± 1 hour).

Intervention Type

Behavioural

Primary outcome(s)

1. Low-density lipoprotein (LDL) cholesterol measured using fasted blood tests at baseline, week 3, 5, 8 and 10
2. Insulin resistance measured using the homeostatic model assessment (HOMA) which utilises fasted glucose and insulin levels, at baseline, week 5 and week 10

Key secondary outcome(s)

1. Weight measured using scales at baseline, week 3, 5, 8 and 10
2. Adiposity measured using the gold standard dual-energy x-ray absorptiometry (DEXA) at baseline and week 10
3. Dietary intake assessed through the use of food diaries at baseline, week 2, 4, 7 and 9, as well as 24 hour dietary recalls at baseline, week 3, 5, 8 and 10
4. Food preferences assessed using questionnaires at baseline, week 3, 5 8 and 10, as well as eye tracking tests at baseline, week 5 and week 10

Completion date

30/06/2023

Eligibility

Key inclusion criteria

Participants must:

1. Be aged 18 – 65 years old
2. Have a BMI ≥ 25 kg/m²
3. Have maintained a stable body weight for the 6 months preceding the study (± 2 kg)
4. Be at increased/moderate/high risk of developing type 2 diabetes, as per the Diabetes UK Diabetes Risk Score (scoring ≥ 7)
5. Be able and willing to give informed oral and informed written consent
6. Complete and meet the defined criteria of pre-study questionnaires
7. Be able and willing to complete daily sleep and food diaries during the study
8. Have an eating period of ≥ 12 hours a day

9. Agree to eat their meals within certain time periods during the day whilst participating in the study

10. Be willing and able to undertake laboratory tests on agreed dates during the study

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

Participants will be excluded if they:

1. Have a specific medical condition as confirmed by pre-study questionnaires – Type 1 or 2 diabetes, Polycystic Ovary Syndrome (PCOS), Schizophrenia, Bipolar illness, Depression or receiving treatment with antipsychotic medication, or have a history of Cardiovascular Disease (myocardial infarction or stroke)
2. Are currently on certain medications: weight loss, glucose- or lipid-lowering medication (e.g. 'Metformin' or 'Statins'), hypnotics or melatonin supplements
3. Are a smoker
4. Are pregnant
5. Have a history of any circadian or sleep disorder as confirmed by pre-study questionnaires
6. Have habitual irregular sleep patterns on more than 2 nights per week (bed time outside 22:00-1:00 h and wake up time outside 6:00-9:00 h) or a sleep duration on more than 2 nights per week < 7 or > 9 hours
7. Have donated over 400 ml of blood in the three months preceding the study
8. Exceed the defined criteria of pre-study questionnaires:
 - 8.1. BMI: < 25kg/m²
 - 8.2. 30 ≤ Horne-Östberg questionnaire ≥70
 - 8.3. Pittsburgh Sleep Quality Index: > 5
 - 8.4. Epworth Sleepiness Scale > 9
9. Have travelled across more than two time zones within a month before or during the study
10. Are currently taking part in a clinical trial or another research study or have taken part within the last 3 months
11. Have participated in rotating or night shift work for more than 6 months prior to or during the study

Date of first enrolment

02/01/2020

Date of final enrolment

31/03/2023

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

University of Surrey

Guildford

Surrey

United Kingdom

GU2 7XH

Sponsor information

Organisation

University of Surrey

ROR

<https://ror.org/00ks66431>

Funder(s)

Funder type

University/education

Funder Name

University of Surrey

Alternative Name(s)

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			26/07/2023	No	No