Study on the safety and efficacy of Aurantii Fructus Immaturus flavonoid extract tablets (Aolanti) in the treatment of functional dyspepsia

Submission date	Recruitment status	Prospectively registered
24/02/2025	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
09/03/2025	Completed	Results
Last Edited	Condition category	[] Individual participant data
07/03/2025	Digestive System	[X] Record updated in last year

Plain English summary of protocol

Background and study aims

This study aims to evaluate the safety and efficacy of Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti) in treating functional dyspepsia (FD) characterized by liver-stomach disharmony, spleen-stomach damp-heat, and food stagnation syndromes. The multicenter clinical trial will explore the therapeutic effects of the drug on FD and provide a foundation for subsequent Phase III clinical trials. This study expects to provide valuable insights into the efficacy and safety of Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti) for FD, supporting its potential as a therapeutic option for patients with specific TCM syndromes.

Who can participate?

Adult participants between the ages of 18 and 65 years old meeting the Western diagnostic criteria for FD and the Traditional Chinese Medicine (TCM) syndrome differentiation criteria were enrolled.

What does the study involve?

Patients will be randomly assigned to either the treatment group, receiving Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti), or the control group, receiving a placebo. The treatment group received 4 tablets of Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti) three times daily, while the control group received 4 placebo tablets under the same regimen, for a total of 4 weeks. The primary efficacy endpoint was the clinical efficacy based on the Western symptom score for FD, while secondary endpoints included clinical efficacy based on TCM syndrome scores and gastric emptying function assessed by the barium strip method.

What are the possible benefits and risks of participating?

Potential benefits included the alleviation of FD symptoms, particularly for patients with liver-stomach disharmony, spleen-stomach damp-heat, and food stagnation syndromes. Potential

risks involved possible adverse drug reactions, although the study results indicated that Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti) were well-tolerated with a favorable safety profile.

Where is the study run from?

The study was conducted across multiple centers in China, including West China Hospital of Sichuan University, Affiliated Hospital of Chengdu University of Traditional Chinese Medicine, The First Affiliated Hospital of Hunan University of Chinese Medicine, Xiangya Hospital of Central South University, and Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine.

When is the study starting and how long is it expected to run for? December 2009 to April 2011

Who is funding the study?

The study was funded by Jiangxi Qingfeng Pharmaceutical Co., Ltd.

Who is the main contact?

Ms Xiaonan Yang, West China Hospital of Sichuan University, yangxxnan@163.com

Contact information

Type(s)

Public

Contact name

Ms Tiancheng Qi

Contact details

10th Floor, Block A, Building 2, Zone 1, Shenzhen Bay Science and Technology Ecological Park, Nanshan District
Shenzhen, Guangdong Province
China
518063
+86-18588477550
qitiancheng@qfyy.com.cn

Type(s)

Scientific

Contact name

Ms Xiaolei Bi

Contact details

9th Floor, Building 3, Kunsha Center, No. 16 Xinyuanli, Chaoyang District Beijing China 100027 +86-13811657829 bixiaolei@qfyy.com.cn

Type(s)

Principal Investigator

Contact name

Ms Xiaonan Yang

Contact details

No. 37, Guoxue Alley, Wuhou District Chengdu, Sichuan Province China 610041 +86-13608000318 yangxxnan@163.com

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Nil known

Study information

Scientific Title

A randomized, double-blind, placebo-controlled, multicenter clinical study on the safety and efficacy of Aurantii Fructus Immaturus flavonoid extract tablets (Aolanti) in the treatment of functional dyspepsia (liver-stomach disharmony, damp-heat in spleen and stomach, and food stagnation syndromes)

Study objectives

Study and evaluate the efficacy and safety of Aurantii Fructus Immaturus flavonoid extract tablets (Aolanti) in treating functional dyspepsia (liver-stomach disharmony, spleen-stomach damp-heat, and food retention syndromes), and explore applicable syndrome types for phase III clinical trials.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 31/12/2009, Drug Clinical Trial Ethics Committee of West China Hospital of Sichuan University (No. 37, Guoxue Alley, Wuhou District, Chengdu, Sichuan Province, 610041, China; +86-28-85422114; hxjj@cd120.com), ref: 2008L04094

Study design

Stratified block-randomized double-blind single-simulation placebo parallel-controlled, multicenter clinical study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment, Safety, Efficacy

Participant information sheet

No participant information sheet available

Health condition(s) or problem(s) studied

The treatment of patients with functional dyspepsia

Interventions

The study adopted a multicenter, randomized, double-blind, placebo-parallel-controlled design, with stratification factors including study center and TCM syndrome type. The ratio of the treatment group to the placebo group was 2:1 within each syndrome subtype. This study divided patients into an experimental group and a control group. The experimental group received Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti) with specifications of 0.29g/tablet (Batch No.: 20090625, provided by Jiangxi Qingfeng Pharmaceutical Co., Ltd.). The dosage was 4 tablets per dose, administered with warm water 30 minutes before meals, three times daily. The control group received a placebo with identical specifications (0.29g/tablet, Batch No.: 20090625, provided by Jiangxi Qingfeng Pharmaceutical Co., Ltd.), following the same administration regimen. The treatment course lasted 28 days, with follow-up visits every 7 days. A post-treatment follow-up period of 28 days was conducted.

Intervention Type

Drug

Pharmaceutical study type(s)

Pharmacodynamic

Phase

Phase II

Drug/device/biological/vaccine name(s)

Aurantii Fructus Immaturus Flavonoid Extract Tablets (Aolanti)

Primary outcome measure

Western Medicine Disease Efficacy: Using the Western medicine primary symptom grading and quantification standards, the four main symptoms are graded and scored, with each symptom

corresponding to a score of 0-3. Assessments are conducted at the screening period, baseline, and on days 7, 14, 21, 28, and 56, with a focus on evaluating the efficacy based on the scores at day 28.

Secondary outcome measures

1. Traditional Chinese Medicine (TCM) Syndrome Efficacy: Using the TCM syndrome grading and quantification standards, patients with syndromes such as Liver-Stomach Disharmony, Spleen-Stomach Dampness-Heat, and Food Retention are scored based on their symptoms. Assessments are conducted at the screening period, baseline, and on days 7, 14, 21, 28, and 56,

with a focus on evaluating the efficacy based on the scores at day 28.

- 2. Gastric Emptying Time: The barium strip method is used to measure the gastric emptying rate at 3.5 hours. Observations and recordings are conducted once before medication (baseline) and once after 28 days of medication.
- 3. Changes in Tongue and Pulse Manifestations: A comprehensive inquiry and physical examination are conducted to observe changes in tongue and pulse before, during, and after treatment. Tongue manifestations are primarily assessed through visual inspection, observing the color of the tongue body and the condition of the tongue coating. Pulse manifestations are primarily assessed through palpation, where the patient's pulse is evaluated to determine its characteristics.

Overall study start date

31/12/2009

Completion date

30/04/2011

Eligibility

Key inclusion criteria

- 1. Sign the informed consent form
- 2. Meet the Western medical diagnostic criteria for functional dyspepsia
- 3. Conform to the Traditional Chinese Medicine (TCM) syndrome diagnostic criteria for epigastric stuffiness (including liver-stomach disharmony, spleen-stomach damp-heat, and food stagnation syndromes)
- 4. Aged 18 to 65 years at the time of enrollment

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

65 Years

Sex

Both

Target number of participants

270

Total final enrolment

265

Key exclusion criteria

- 1. Individuals with organic diseases such as esophagitis, atrophic gastritis, gastric and duodenal ulcers, erosions, tumors, etc.
- 2. Individuals with organic diseases of the liver, gallbladder, or pancreas
- 3. Individuals with diabetes, kidney disease, connective tissue disease, or psychiatric disorders
- 4. Individuals with severe primary diseases of the heart, brain, lungs, kidneys, hematopoietic system, or endocrine system
- 5. Individuals with a history of abdominal surgery
- 6. Pregnant women, those with pregnancy intentions, or women who are breastfeeding
- 7. Individuals with mental or legal disabilities
- 8. Individuals who have participated in other drug clinical trials within the 3 months prior to this trial

Date of first enrolment

01/03/2010

Date of final enrolment

08/12/2010

Locations

Countries of recruitment

China

Study participating centre

West China Hospital of Sichuan University

No. 37, Guoxue Alley, Wuhou District Chengdu, Sichuan Province China 610041

Study participating centre

Affiliated Hospital of Chengdu University of Traditional Chinese Medicine

No. 39, Shierqiao Road, Jinniu District Chengdu, Sichuan Province China 610072

The First Affiliated Hospital of Hunan University of Chinese Medicine

No. 95, Shaoshan Middle Road, Furong District Changsha, Hunan Province China 410007

Study participating centre Xiangya Hospital of Central South University

No. 87, Xiangya Road, Kaifu District Changsha, Hunan Province China 410008

Study participating centre

Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine

No. 445, Bayi Avenue, Xihu District Nanchang, Jiangxi Province China 330006

Sponsor information

Organisation

Jiangxi Qingfeng Pharmaceutical Co., Ltd.

Sponsor details

Shahe Industrial Park, Ganzhou Jiangxi Province China 341000 +86 0797-5560800 yxpv@qfyy.com.cn

Sponsor type

Industry

Website

https://www.qfyy.com.cn/

Funder(s)

Funder type

Industry

Funder Name

Jiangxi Qingfeng Pharmaceutical Co., Ltd.

Results and Publications

Publication and dissemination plan

Planned publication in a peer-reviewed journal

Intention to publish date

31/12/2025

Individual participant data (IPD) sharing plan

The aggregated research data will be stored by the sponsor as required. The original research data will be retained at each research center in accordance with regulations. Interested parties may submit a request to the sponsor as needed, and the sponsor will provide relevant content based on actual circumstances. (Contact: Ms Qi, qitiancheng@qfyy.com.cn)

IPD sharing plan summary

Available on request