# Ultrasound contrast agents to facilitate sonothrombolysis in patients with acute myocardial infarction

Submission date	Recruitment status	Prospectively registered
20/12/2005	No longer recruiting	[X] Protocol
Registration date	Overall study status	Statistical analysis plan
20/12/2005	Completed	[X] Results
Last Edited	Condition category	Individual participant data
20/08/2021	Circulatory System	

# Plain English summary of protocol

Not provided at time of registration

## Contact information

## Type(s)

Scientific

#### Contact name

Dr O Kamp

#### Contact details

VU University Medical Centre Department of Cardiology, 6D-120 P.O. Box 7057 Amsterdam Netherlands 1007 MB +31 (0)20 444 2244 o.kamp@vumc.nl

## Additional identifiers

**EudraCT/CTIS** number

**IRAS** number

ClinicalTrials.gov number

Secondary identifying numbers

# Study information

#### Scientific Title

Ultrasound contrast agents to facilitate sonothrombolysis in patients with acute myocardial infarction

#### Acronym

**ULYSIS** 

#### **Study objectives**

The optimal treatment strategy in patients with Acute Myocardial Infarction (AMI) is immediate restoration of coronary blood flow. Although thrombolytic therapy is the most widely used therapy, Percutaneous Coronary Intervention (PCI) is the treatment of choice in AMI patients, however, its widespread use is hampered by limited availability of specialised facilities and trained staff. There is a need for simpler and low-risk methods for effective recanalisation of thrombosed arteries that can be initiated early in the disease process.

Recently, the application of ultrasound in combination with thrombolytic agents was found to enhance thrombus dissolution in vitro and in vivo. In vivo studies using thrombo-occlusive canine and rabbit models demonstrated that Ultrasound Contrast Agents (UCAs) enhance this thrombus dissolving effect of ultrasound, resulting in higher recanalisation rates of occluded arteries.

#### Hypothesis:

Under influence of ultrasound, UCAs enhance dissolution of thrombus in patients with acute ST-elevation myocardial infarction premedicated with aspirin and clopidogrel.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised, placebo controlled, parallel group, single blinded multicentre trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Not specified

## Study type(s)

**Treatment** 

#### Participant information sheet

#### Health condition(s) or problem(s) studied

Acute myocardial infarction

#### **Interventions**

After having received 150 - 325 mg aspirin and a loading dose of 300 mg clopidogrel, patients will be randomised to:

- 1. Ultrasound application with infusion of an ultrasound contrast agent, or
- 2. Control (infusion of saline without ultrasound application).

Immediately after ultrasound application, catheterisation will be performed.

#### **Intervention Type**

Drug

#### Phase

**Not Specified** 

#### Drug/device/biological/vaccine name(s)

Aspirin and clopidogrel

#### Primary outcome measure

Comparison of the UCA-group and the control group with respect to:

- 1. Patency of the culprit coronary artery
- 2. Thrombolysis In Myocardial Infarction (TIMI)-flow
- 3. Corrected TIMI frame count
- 4. Myocardial blush grade

#### Secondary outcome measures

- 1. ST-segment resolution
- 2. Release of cardiac enzymes
- 3. Echocardiographic wall motion score index
- 4. Safety

#### Overall study start date

01/10/2005

#### Completion date

01/12/2006

# Eligibility

#### Key inclusion criteria

- 1. Age 18 to 80 years
- 2. Diagnosed with acute myocardial infarction according to the criteria of the American College of Cardiology
- 3. Informed consent

#### Participant type(s)

#### **Patient**

#### Age group

Adult

## Lower age limit

18 Years

#### Sex

**Not Specified** 

#### Target number of participants

60

### Key exclusion criteria

- 1. Previous myocardial infarction
- 2. Clinical instability
- 3. Pregnancy/breast feeding
- 4. Known pulmonary hypertension
- 5. Known allergy to Optison
- 6. Any reason judged by the investigators to hamper inclusion

#### Date of first enrolment

01/10/2005

#### Date of final enrolment

01/12/2006

## Locations

#### Countries of recruitment

Netherlands

# Study participating centre VU University Medical Centre

Amsterdam Netherlands 1007 MB

# Sponsor information

#### Organisation

VU University Medical Centre (The Netherlands)

## Sponsor details

Van der Boechorststraat 7 Amsterdam Netherlands 1081 BT

## Sponsor type

University/education

#### Website

http://www.vumc.nl/english/

#### **ROR**

https://ror.org/00q6h8f30

# Funder(s)

## Funder type

Not defined

#### Funder Name

Not provided at time of registration

## **Results and Publications**

## Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article		08/03/2011	20/08/2021	Yes	No
Results article		01/02/2012	20/08/2021	Yes	No