

Pelvic (floor) Reminders (to) Explore Perinatal (women's) Acceptability (of) Reminders (to) Exercise (PREPARE) study

Submission date 26/08/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
Registration date 27/09/2023	Overall study status Completed	<input checked="" type="checkbox"/> Protocol <input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 09/04/2025	Condition category Urological and Genital Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Pelvic floor muscle dysfunction (PFMD) is a condition that affects up to one in three women at some point in their lifetime and can be commonly traced back to the perinatal period due to the physiological changes associated with pregnancy and childbirth. The most prevalent symptom of PFMD is stress urinary incontinence (SUI), defined as the involuntary loss of urine on effort or physical exertion (e.g., sporting activities), or on sneezing or coughing. The social, emotional, economic, and environmental impact of SUI is well documented and includes social isolation, difficulty returning to work, impaired sexual function and prevention of vigorous exercise. Supervised pelvic floor muscle training (PFMT) is the gold standard of treatment for symptoms of SUI. Pelvic floor muscle training is defined as exercise to improve pelvic floor muscle strength, endurance, power, relaxation or a combination of these parameters. Women are less likely to develop urinary incontinence during pregnancy or in the early post-natal period if they regularly complete PFMT. PFMT involves women completing supervised PFMT three times a day for a minimum of three months for it to be effective at reducing symptoms of PFMD; however, clinically, it is recognised women do not regularly adhere to this exercise prescription. Health apps commonly use reminders for long-term conditions to promote self-management and have been proven effective in other long-term conditions. The use of digital technology, such as mobile apps, in maternity care can personalise the treatment individual women receive thus improving healthcare delivery and in particular the management of PFMD. Personalisation of PFMT may be enhanced by providing digital 'nudges' as reminders and encouragement to enhance adherence to exercises.

Who can participate?

Women between the ages of 18 and 45 years who have a viable pregnancy and have given birth once before

What does the study involve?

The feasibility trial will determine whether a future randomised controlled trial can be conducted. The feasibility trial will compare an intervention and a control group. The intervention group will receive a PFMT mobile app and digital nudges. The control group will

receive a leaflet on the role of the pelvic floor and how to complete PFMT developed by a UK-based professional network of clinical specialist physiotherapists. Perinatal women will have three telephone follow-ups lasting no longer than 30 minutes. The timing of these follow-ups will be at 9, 21 and 31 weeks. These follow-ups correspond to the late antenatal stage of gestation (36 weeks), around 1 month postnatal and 4 months postnatal.

What are the possible benefits and risks of participating?

The immediate advantage to participants is support with pelvic floor muscle training during pregnancy. Whilst the researchers do not anticipate any risks to participants there is a very small risk of emotional distress talking about pelvic floor-related symptoms. For those in the intervention group, there is a small but possible chance that participants find digital reminders a burden. There is a small but possible risk participants may find the statements in the digital nudges hard to understand. Participants randomised into a group they are not happy with may experience negative feelings.

Where is the study run from?

Bournemouth University (UK)

When is the study starting and how long is it expected to run for?

January 2023 to April 2025

Who is funding the study?

1. Bournemouth University (UK)
2. University Hospitals Dorset (UK)
3. National Institute for Health Research (NIHR) Applied Research Collaboration (ARC) Wessex (UK)

Who is the main contact?

Rosie Harper, rharper1@bournemouth.ac.uk

Contact information

Type(s)

Public

Contact name

Miss Rosie Harper

ORCID ID

<https://orcid.org/0000-0003-3414-7580>

Contact details

Faculty of Health and Sports Sciences

Bournemouth Gateway Building

12 St Paul's Ln

Bournemouth

United Kingdom

BH8 8GP

+44 (0)1202 969696

rharper1@bournemouth.ac.uk

Type(s)

Scientific

Contact name

Miss Rosie Harper

Contact details

Faculty of Health and Sports Sciences
Bournemouth Gateway Building
12 St Paul's Ln
Bournemouth
United Kingdom
BH8 8GP
+44 (0)1202 969696
rharper1@bournemouth.ac.uk

Type(s)

Principal investigator

Contact name

Prof Carol Clark

ORCID ID

<https://orcid.org/0000-0002-9296-7141>

Contact details

Faculty of Health and Sports Sciences
Bournemouth Gateway Building
12 St Paul's Ln
Bournemouth
United Kingdom
BH8 8GP
+44 (0)1202 969696
cclark@bournemouth.ac.uk

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Integrated Research Application System (IRAS)

328225

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

IRAS 328225, CPMS 58654

Study information

Scientific Title

Pelvic (floor) Reminders (to) Explore Perinatal (women's) Acceptability (of) Reminders (to) Exercise (PREPARE)

Acronym

PREPARE

Study objectives

The principal research objective is to determine the feasibility of undertaking a randomized controlled trial that explores whether a physiotherapy-led digital intervention can increase pelvic floor muscle training adherence in women.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. Approved 19/12/2023, HRA REC South Central (South Central – Hampshire A, Southampton, SO17 1BJ, United Kingdom; +44 (0)207 104 8120, +44 (0)207 104 8210, +44 (0)207 104 8290; hampshirea.rec@hra.nhs.uk), ref: 23/SC/0407
2. Approved 12/01/2024, Bournemouth University (Faculty of Health and Sports Sciences, Bournemouth, BH8 8GP, United Kingdom; +44 (0)1202 524111; researchethics@bournemouth.ac.uk), ref: 52246

Study design

Single-centre interventional mixed methods feasibility study

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Pelvic floor dysfunction

Interventions

Participants will be randomised between 26-28 weeks of pregnancy by assigning random numbers to each subject using permuted block randomisation in REDCap™ software. Participants will be randomised into one of two arms of the study: the intervention group or the control group. The researchers are unable to blind the participants in the study. It was not possible to blind data collectors in the study. The control group will receive a leaflet on the role of the pelvic floor and how to complete PFMT developed by a UK-based professional network of clinical specialist physiotherapists.

The intervention consists of 31 targeted digital PFMT 'nudges' and a pelvic floor muscle training mobile app. The digital nudges were co-designed as part of a larger research study to support perinatal women to complete regular pelvic floor muscle training. The total number of digital nudges is 31 to account for the number of weeks between 26 weeks of pregnancy and 4 months postnatal. The nudges target different stages of the perinatal period and have been divided into the following categories:

1. Late antenatal
2. Early postnatal
3. Late postnatal

The nudges will be in the form of push notifications. A digital nudge will be sent once a week to each participant. Women will receive the first digital nudge between 26-28 weeks of pregnancy soon after they have been enrolled in the study. Each digital nudge is unique and the last digital nudge will be sent around 4 months post-natal.

The control group will receive a leaflet on the role of the pelvic floor and how to complete PFMT developed by a UK-based professional network of clinical specialist physiotherapists.

Intervention Type

Behavioural

Primary outcome(s)

Several outcome measures will be trialled in the study to determine whether the study design and intervention can be taken forward for an RCT. Primary outcome measures include:

1. Quality of life measured using EQ5D5L at baseline, 9 weeks, 21 weeks and 31 weeks
2. Severity of urinary incontinence symptoms measured using International Consultation of Incontinence Questionnaire of Urinary Incontinence- Short Form (ICIQ-UI) at baseline, 9 weeks, 21 weeks and 31 weeks
3. Self-efficacy around pelvic floor muscle training measured using the Broome pelvic muscle self-efficacy scale at baseline and 31 weeks
4. Adherence to pelvic floor muscle training measured using in-app adherence data that will be collected weekly throughout the study
5. Adherence to pelvic floor muscle training measured using online and paper diaries at 9 weeks, 21 weeks and 31 weeks
6. Participant engagement with the intervention measured using in-app algorithms e.g. interaction response time and will be collected continuously throughout the study

Key secondary outcome(s)

Secondary outcome measures will be used to add greater depth to the quantitative data collected in the study to explore participants' acceptability of the intervention.

The secondary outcome measures include:

1. Participant's views on the acceptability of the digital nudges and study processes will be measured using semi-structured interviews at just after 31 weeks
2. Participant's acceptability of the intervention will be captured using one open-ended questions 'which nudges best supported you in your training and why' at 9 weeks, 21 weeks and 31 weeks

Completion date

30/04/2025

Eligibility

Key inclusion criteria

Primiparous women over the age of 18 years with a viable pregnancy and the ability to send and receive e-mail

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

45 years

Sex

Female

Total final enrolment

44

Key exclusion criteria

1. Under the age of 18 years
2. Haematuria
3. Difficulty passing urine or bladder emptying difficulties
4. Present malignancy of the pelvic area
5. A neurological disease that affects the urinary system
6. Pyelonephritis
7. Severe comorbidities in pregnancy (including placenta previa, threatened premature labour, pregnancy-induced hypertension)
8. Hyperactivity of the pelvic floor
9. Active urinary tract infection
10. History of stroke, diabetes or gestational diabetes
11. Use of another PFMT mobile app
12. Cannot read or understand written English

Date of first enrolment

23/02/2024

Date of final enrolment

23/05/2024

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

University Hospitals Dorset NHS Foundation Trust
Management Offices
Poole Hospital
Longfleet Road
Poole
United Kingdom
BH15 2JB

Sponsor information

Organisation

Bournemouth University

ROR

<https://ror.org/05wwcw481>

Funder(s)

Funder type

University/education

Funder Name

Bournemouth University

Alternative Name(s)

Bournemouth Municipal College, Bournemouth College of Technology, Dorset Institute of Higher Education, Bournemouth Polytechnic, BU, DIHE

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Funder Name

National Institute for Health and Care Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

University Hospitals Dorset

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Rosie Harper (rharper1@bournemouth.ac.uk). Consent from participants will be required and obtained and participants will be pseudo-anonymised using a unique study identity code. All other details will be added to the study record at a later date.

IPD sharing plan summary

Stored in non-publicly available repository, Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol (preprint)		19/01/2024	13/05/2024	No	No