

Development of endoscopic Optical Coherence Tomography for Laryngeal cancer screening

Submission date	Recruitment status	<input type="checkbox"/> Prospectively registered
16/08/2011	No longer recruiting	<input type="checkbox"/> Protocol
Registration date	Overall study status	<input type="checkbox"/> Statistical analysis plan
02/03/2012	Completed	<input checked="" type="checkbox"/> Results
Last Edited	Condition category	<input type="checkbox"/> Individual participant data
12/12/2017	Cancer	

Plain English summary of protocol

Background and study aims

Larynx (voicebox) cancer is one of the most common head and neck cancers. The main risk factors are smoking and drinking alcohol. Early larynx cancer can be treated by laser, which has a cure rate of over 90% and avoids the side effects of radio/chemotherapy. Later stage larynx cancer is hugely expensive to treat and wasteful of healthcare resources, often requiring mutilating treatment resulting in loss of ability to speak or swallow. Optical Coherence Tomography (OCT) is a relatively new technology where light is used to scan a tissue's surface and a few millimetres deep into it, obtaining accurate information about the tissue's characteristics, without the need to physically cut into tissue in order to microscopically examine if it is normal, precancerous or cancerous. We aim to test whether new OCT probes can distinguish normal from unhealthy larynx tissue and determine whether it is acceptable for clinicians and patients.

Who can participate?

Patients aged over 18 with suspicious larynx appearances on endoscopy.

What does the study involve?

Participants will have endoscopic OCT images taken before going on to have biopsies (samples) of the larynx tissue taken under anaesthetic. One pathologist will compare the OCT results with the standard biopsy results to determine the effectiveness of OCT in distinguishing cancerous from noncancerous tissue. Patient acceptability of the process will be established through a patient experience questionnaire.

What are the possible benefits and risks of participating?

Following this study we aim to use the results to develop a non-invasive, patient-friendly and safe laryngeal cancer screening tool. With this we aim to increase the early detection and treatment of larynx cancer.

Where is the study run from?

North West London Hospitals NHS Trust (UK).

When is the study starting and how long is it expected to run for?
October 2011 to February 2012.

Who is funding the study?
National Institute for Health Research (UK).

Who is the main contact?
Taran Tatla
ttatla@nhs.net

Contact information

Type(s)
Scientific

Contact name
Mr Taran Tatla

Contact details
North West London Hospitals NHS Trust
Watford Road
Harrow
London
United Kingdom
HA1 3UJ
-
ttatla@nhs.net

Additional identifiers

Protocol serial number
8939

Study information

Scientific Title
Development of endoscopic optical coherence tomography for laryngeal cancer screening: a non-randomised trial

Acronym
OCTiLarynx

Study objectives
OCTiLarynx (Optical Coherence Tomography in Larynx). This 'proof of concept' project has a primary objective to determine if recent and continuing advances in the technology of Optical Coherence Tomography (OCT) can result in a clinically useful out-patient based diagnostic tool for endoscopic laryngeal disease screening.

We aim to validate the clinical efficacy of novel endoscopic OCT imaging probes for distinguishing normal from unhealthy laryngeal mucosa (inflamed, dysplastic, malignant) and determine clinician/patient acceptability for the same.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Central London Research Ethics Committee, 28/07/2010, ref: 10/H0718/55

Study design

Non randomised trial

Primary study design

Interventional

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Laryngeal cancer screening

Interventions

Clinical recruitment of up to 40 patients for in vivo study shall occur initially at a single site (Northwick Park ENT-Head & Neck Department). Patients attending the outpatient service for secondary / tertiary care consult in relation to prolonged voice change and noted to have a suspect laryngeal mucosal lesion shall be recruited prospectively for the study. Following informed consent, these patients shall have standard endoscopic digital photographic images and endoscopic OCT images of the lesions taken under topical local anaesthesia (LA) in the outpatient department. As is standard practice presently, all such patients shall then proceed on to direct biopsies of the laryngeal lesions under general anaesthesia to allow gold standard histological diagnosis with routine Haematoxylin and Eosin stains of wax-embedded tissue sections.

A single pathologist trained in OCT tissue imaging techniques shall make a blinded comparison, comparing and correlating the OCT diagnostic modality with the standard biopsy histological technique for distinguishing laryngeal mucosal disease category. Preliminary statistical analysis will determine the efficacy and repeatability of the OCT tool in distinguishing cancer from non-cancer. Patient acceptability of the process will be established through a patient experience questionnaire and clinician feedback shall be sought on practicality and ease of use of the endoscopic OCT tool.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Characterisation & correlation of mucosal pathology (OCT images vs Histopathology) following direct microlaryngoscopy and biopsy of laryngeal lesion under general anaesthesia (GA)

Key secondary outcome(s)

Patient Experience Questionnaire at initial out-patient visit , immediately following nasendoscopy examination and image capture under LA

Completion date

29/02/2012

Eligibility

Key inclusion criteria

1. All patients over the age of 18 years (no upper age limit)
2. Male or female gender
3. Patients able to provide informed consent to OCT-adapted flexible nasendoscope examination under topical local anaesthesia and completing a patient experience questionnaire provided at the end of procedure
4. Smokers and non-smokers

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. All patients under the age of 18 years
2. All patients from whom informed consent has not been obtained prior to tissue resection or OCT-adapted flexible nasendoscopy respectively

Date of first enrolment

03/10/2011

Date of final enrolment

29/02/2012

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

North West London Hospitals NHS Trust
London
United Kingdom
HA1 3UJ

Sponsor information

Organisation

North West London Hospitals NHS Trust (UK)

ROR

<https://ror.org/04cntmc13>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<u>Results article</u>	results	01/12/2012		Yes	No
<u>Participant information sheet</u>	Participant information sheet	11/11/2025	11/11/2025	No	Yes