Does applying limb compression before long saphenous vein stripping result in a better outcome than applying it after stripping?

Submission date	Recruitment status	Prospectively registered
30/09/2005	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
30/09/2005	Completed	Results
Last Edited	Condition category	Individual participant data
14/06/2017	Surgery	[] Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Mr Sanjeev Sarin

Contact details

Surgery
Watford General Hospital
Vicarage Road
Watford
United Kingdom
WD18 0HB
+44 1923 244366
sanjeev.sarin@whht.nhs.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

Study information

Scientific Title

Does applying limb compression before long saphenous vein stripping result in a better outcome than applying it after stripping?

Study objectives

To find out whether applying limb compression before the long saphenous vein is stripped results in a better postoperative outcome than applying it at the end of the procedure.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Double-blinded block randomisation pilot study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Surgery: Varicose vein

Interventions

Limb compression before the long saphenous vein is stripped compared to after.

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome measure

Post-operative outcome.

Secondary outcome measures

Not provided at time of registration

Overall study start date

17/05/2004

Completion date

01/04/2006

Eligibility

Key inclusion criteria

25 patients requiring varicose vein surgery

Participant type(s)

Patient

Age group

Not Specified

Sex

Not Specified

Target number of participants

25

Key exclusion criteria

Patients requiring saphenopopliteal ligation as well

Date of first enrolment

17/05/2004

Date of final enrolment

01/04/2006

Locations

Countries of recruitment

England

United Kingdom

Study participating centre Watford General Hospital

Watford United Kingdom WD18 0HB

Sponsor information

Organisation

Department of Health

Sponsor details

Richmond House 79 Whitehall London United Kingdom SW1A 2NL +44 (0)20 7307 2622 dhmail@doh.gsi.org.uk

Sponsor type

Government

Website

http://www.dh.gov.uk/Home/fs/en

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

West Hertfordshire Hospitals NHS Trust (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration