

Does applying limb compression before long saphenous vein stripping result in a better outcome than applying it after stripping?

Submission date 30/09/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 30/09/2005	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 14/06/2017	Condition category Surgery	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr Sanjeev Sarin

Contact details
Surgery
Watford General Hospital
Vicarage Road
Watford
United Kingdom
WD18 0HB
+44 1923 244366
sanjeev.sarin@whht.nhs.uk

Additional identifiers

Protocol serial number
N0143157279

Study information

Scientific Title

Does applying limb compression before long saphenous vein stripping result in a better outcome than applying it after stripping?

Study objectives

To find out whether applying limb compression before the long saphenous vein is stripped results in a better postoperative outcome than applying it at the end of the procedure.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Double-blinded block randomisation pilot study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Surgery: Varicose vein

Interventions

Limb compression before the long saphenous vein is stripped compared to after.

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome(s)

Post-operative outcome.

Key secondary outcome(s)

Not provided at time of registration

Completion date

01/04/2006

Eligibility**Key inclusion criteria**

25 patients requiring varicose vein surgery

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Patients requiring saphenopopliteal ligation as well

Date of first enrolment

17/05/2004

Date of final enrolment

01/04/2006

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Watford General Hospital

Watford

United Kingdom

WD18 0HB

Sponsor information**Organisation**

Department of Health

Funder(s)**Funder type**

Hospital/treatment centre

Funder Name

West Hertfordshire Hospitals NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration