

# Dietary oligofructose in the prevention of antibiotic associated diarrhoea in elderly patients

<b>Submission date</b> 12/09/2003	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 12/09/2003	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 06/01/2009	<b>Condition category</b> Digestive System	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Stephen Lewis

**Contact details**  
Box No 201A  
Department of Gastroenterology  
Addenbrooke's NHS Trust  
Cambridge  
United Kingdom  
CB2 2QQ  
+44 (0)7669 008 863

## Additional identifiers

**Protocol serial number**  
N0544103828

## Study information

**Scientific Title**

**Study objectives**

The purpose of this study is to see if the incidence of antibiotic related diarrhoea in elderly patients can be reduced by increasing colonic bifidobacteria concentrations with the concomitant prescription of oligofructose. Patients for the study will be identified by asking the Medical and Department for the Elderly teams if they had admitted any suitable patients. A brief clinical and drug history would be taken.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Randomised controlled trial

**Primary study design**

Interventional

**Study type(s)**

Not Specified

**Health condition(s) or problem(s) studied**

Digestive System: Diarrhoea

**Interventions**

The volunteers would then be randomly allocated to one of two groups.

Group A will take a placebo (10 g/day of sucrose powder) or group B 10 g/day of oligofructose. The trial powders will be administered by the ward nursing staff along with the volunteers' normal medication. Volunteers will take the trial powders for the length of time they are taking antibiotics and for a further 7 days after stopping antibiotics. Volunteers will be assessed continuously by recording on a stool form using a four point scale (1 hard, 2 lumpy, 3 mushy and 4 loose) and interdefecatory intervals. Both give an indication of intestinal transit rate. Recordings will be done by the patient or nursing staff as appropriate. The volunteers will be followed up for 7 days after stopping the oligofructose or placebo. The volunteer will have further stool samples sent to be analysed for the presence of Clostridium difficile toxin if they develop diarrhoea during the study. The study's end point is the occurrence of diarrhoea due to Clostridium difficile.

**Intervention Type**

Other

**Phase**

Not Specified

**Primary outcome(s)**

Not provided at time of registration

**Key secondary outcome(s)**

Not provided at time of registration

**Completion date**

15/06/2003

## Eligibility

**Key inclusion criteria**

250 volunteers >65 years old.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Senior

**Sex**

Not Specified

**Key exclusion criteria**

Not provided at time of registration

**Date of first enrolment**

16/06/2000

**Date of final enrolment**

15/06/2003

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre**

Box No 201A

Cambridge

United Kingdom

CB2 2QQ

## Sponsor information

**Organisation**

Department of Health (UK)

**Funder(s)****Funder type**

Government

**Funder Name**

Cambridge Consortium - Addenbrooke's (UK)

**Results and Publications****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	15/02/2005		Yes	No