

Waist circumference as an estimator of abdominal wall depth

Submission date 08/01/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 02/02/2023	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 26/08/2025	Condition category Other	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

This study seeks to assess whether waist circumference can estimate the abdominal wall depth to a reliable degree of accuracy so as to be incorporated in a nomogram that would categorise suggested Veress needle abdominal insertion depth according to waist circumference range. This would be helpful in guiding initial Veress entry into the abdominal cavity during laparoscopic (keyhole) or robotic surgery where entry is achieved blindly without standardised depth advice. The research question is therefore 'Can waist circumference estimate abdominal wall depth?' Due to the urgent need for safety mechanisms to be incorporated in the conduct of laparoscopic abdominal wall injury, this study attempts to provide some guidance into how deep the Veress needle should be inserted when a Veress needle is being used as the first entry abdominal device and insufflator. The study aims to improve safety in laparoscopic entry where a Veress needle is used by providing insertion depth guidance. It is aimed to reduce the number of failed entries or complicated entries. The traditional Body Mass Index (BMI) will also be taken to see if it makes a better correlation than abdominal girth.

Who can participate

Any adult patient undergoing non-urgent scheduled CT or MRI abdominal imaging

What does the study involve?

The abdominal wall depth will be measured directly from CT or MRI scans of the waist, whilst patient abdominal circumference, weight and height measurements will be physically obtained.

What are the possible benefits and risks of participating?

The possible benefits of participating are that the research team will assess the validity of the relationship between waist circumference and abdominal wall depth for the first time so this can be taken into account whenever patients are subjected to minimally invasive surgery where entry is likely to be initially accomplished through the use of the Veress needle. There is minimal risk to participation, if any, there may be an inconvenience in taking the physical measurements. Weight and height are routinely taken, however, additionally taking the waist circumference is not envisaged to create significant delay.

Where is the study run from?

University Hospitals Bristol and Weston NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?

April 2022 to July 2024

Who is funding the study?

This study is part of a PhD project funded by UK Research and Innovation (UKRI) ESRC Centres for Doctoral Training Digital Health & Care (UK)

Who is the main contact?

Chimwemwe Miti (Principal researcher and PhD student), c.miti@bristol.ac.uk (UK)

Contact information

Type(s)

Principal investigator

Contact name

Miss Chimwemwe Miti

ORCID ID

<https://orcid.org/0000-0001-8309-269X>

Contact details

EPSRC Centre for Doctoral Training (CDT) in Digital Health & Care

School of Computer Science

Electrical and Electronic Engineering and Engineering Maths

1 Cathedral Square

Trinity Street

Bristol

United Kingdom

BS1 5DD

+44 (0)1782730733

zb20006@bristol.ac.uk

Type(s)

Scientific

Contact name

Dr Hermes Gadelha- Bloomfield

ORCID ID

<https://orcid.org/0000-0001-8053-9249>

Contact details

Ada Lovelace Building

Department of Engineering Mathematics

University Walk

Bristol

United Kingdom

BS8 1TW
+44 (0)117 4282489
hermes.gadelha@bristol.ac.uk

Type(s)

Public

Contact name

Dr Lynn Armstrong

Contact details

St Michael's Hospital
University Hospitals Bristol and Weston NHS Foundation Trust
Bristol
United Kingdom
BS2 8EG
None provided
lynne.armstrong@uhbw.nhs.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

321171

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

IRAS 321171, CPMS 55619

Study information

Scientific Title

Investigating whether the abdominal waist circumference or abdominal girth can be employed as an accurate estimate of abdominal wall depth as measured from the base of the umbilicus

Study objectives

Measured waist circumference is a better estimator than Body Mass Index (BMI) of the abdominal wall depth

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 21/07/2023, London - Harrow Research Ethics Committee (Health Research Authority Bristol HRA Centre, Ground Floor, Temple Quay House, 2 The Square, Bristol, BS1 6PN, United Kingdom; +44 (0)2071048289; harrow.rec@hra.nhs.uk), ref: 23/PR/0600

Study design

Observational cross-sectional study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Assessing the correlation between waist circumference and abdominal wall depth

Interventions

For eligible participants undergoing cross-sectional imaging (CT or MRI scans) as part of their routine diagnostic workup, their waist circumference (additional to height and weight which are already routinely collected before any CT or MRI scan) will be measured. The principal investigator who is a PhD student and a medical doctor trained in taking patient physical parameters will measure the waist circumference, height and weight. The measurements of physical parameters are undertaken face-to-face in the imaging department where the patient will be attending their scheduled scan. These data will be analysed mathematically against the abdominal wall depth taken from the base of the umbilicus to the peritoneum and which will be measured directly from an appropriate slice of the cross-sectional image series.

Intervention Type

Other

Primary outcome(s)

1. Abdominal wall depth from the centre of the umbilicus to the peritoneal edge in millimetres (mm) measured using cross-sectional imaging (CT or MRI scans) at one timepoint
2. Waist circumference measured in a complete circle passing through the centre of the umbilicus in millimetres (mm) using a measuring tape at one timepoint
3. Height and weight measured using routine procedures in centimetres (cm) and kilograms (kg), respectively, at one timepoint

Key secondary outcome(s)

Performance of the relationship between waist circumference measurements and abdominal wall depth and measured BMI to estimate abdominal wall depth at a single time point

Completion date

21/07/2024

Eligibility

Key inclusion criteria

1. Adult patients aged 18 years old and over
2. Undergoing routine scheduled cross-sectional imaging that includes their abdomen
3. Mobile patients
4. Able to consent and volunteer their participation in the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

386

Key exclusion criteria

1. Patients with uncorrected abdominal wall hernias or abdominal wall defects
2. Patients undergoing emergency CT scans requested from the emergency department

Date of first enrolment

31/07/2023

Date of final enrolment

24/06/2024

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

University Hospitals Bristol & Weston

Radiology Department

Bristol Royal Infirmary

Marlborough Street

Bristol

United Kingdom

BS2 8HW

Sponsor information

Organisation

University Hospitals Bristol and Weston NHS Foundation Trust

Funder(s)

Funder type

Government

Funder Name

UK Research and Innovation

Alternative Name(s)

UKRI

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and /or analysed during the current study will be stored in a publicly available repository. The Research Data Storage Facility, Digital Health Doc (digitaldoc.info) will be the most robust long-term storage facility for my project data so that the data is accessible by designated researchers should I not be present and when the project ends and I leave the University. The type of data stored will be metric measurements of abdominal wall depth, height, weight, waist circumference, and Body Mass Index. Consent from participants was required and obtained. Data anonymization was as per protocol. There are no ethical or legal restrictions foreseen because only measurements being taken will be stored, the scan images will not be stored.

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results			26/08/2025	No	No
Participant information sheet	version 1.6	02/07/2021	16/01/2023	No	Yes
Protocol file	version 1.0	31/12/2022	16/01/2023	No	No

