

Invisalign - a concept for the prevention of white spot lesions in teenaged orthodontic patients?

Submission date 03/09/2014	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 16/10/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 16/10/2014	Condition category Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Wire dental braces (fixed vestibular multi-bracket appliances) are traditionally considered to be the best treatment (gold standard) for correcting misaligned teeth (dental malocclusion) in teenage patients. However, patients wearing wire braces are 5 times more likely to develop dental cavities (caries) and white spot lesions (often an early sign of tooth decay) than people who don't. Such damage to the teeth cannot be reversed. Invisalign is an alternative to the wire brace. It is made out of plastic, using transparent aligners to gradually adjust the affected teeth. It is often used by people who have misaligned teeth but want a more attractive solution to wearing the traditional brace. Here, we want to find out whether wearing Invisalign rather than wire dental braces will result in fewer or more white spot lesions after 6 months and then after 12 months of therapy.

Who can participate?

Patients aged between 13-19 years needing dental (orthodontic) treatment to correct misaligned teeth.

What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 are treated with a traditional multi-bracket appliance for orthodontic treatment. Those in group 2 receive Invisalign therapy. All participants are examined for white spot lesions at the start of the study, after 6 months of orthodontic treatment, and then after 12 months of treatment.

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

Department of Orthodontics, University Hospital Freiburg (Germany)

When is the study starting and how long is it expected to run for?

November 2014 to October 2016

Who is funding the study?
International Align Research Award of the Align Technology, North America (USA)

Who is the main contact?
Professor Britta A. Jung

Contact information

Type(s)
Scientific

Contact name
Prof Britta A. Jung

Contact details
Head of the Department of Orthodontics
Department of Orthodontics
University Hospital Freiburg, Germany
Hugstetter Str. 55
Freiburg, Germany
Germany
79106

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N/A

Study information

Scientific Title
Invisalign - a concept for the prevention of white spot lesions in teenaged orthodontic patients?
A prospective controlled clinical trial

Study objectives
The aim of this study is to investigate the incidence of white spot lesions during orthodontic treatment with Invisalign and multi-bracket appliance treatment (the gold standard) in teenaged patients.
The null hypothesis of no difference in the incidence of white spot lesions between teeth undergoing treatment with fixed braces and Invisalign therapy will be investigated. The investigation will be designed as a prospective, bicenter controlled clinical study.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of Albert-Ludwigs-University of Freiburg, Germany (Ref. No: 406/14 (MPG§23))

Study design

A prospective controlled clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Other

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Malocclusion

Interventions

Two groups of patients (male and female, aged 13-19 years) will be randomly allocated into one of two arms:

1. Group 1 will consist of 30 patients. All of these patients receive a traditional multi-bracket appliance for orthodontic treatment.
2. Group 2 will consist of 30 patients. All of these patients receive Invisalign therapy for orthodontic treatment.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

1. Macroscopic characterization (presence, frequency, and severity) of white spot lesions at baseline, after 6 months, and after 12 months of orthodontic treatment. The severity of disease will be rated according to Gorelick's classification:

Grade 0: no visible white spot lesions on clinical investigation

Grade 1: minor white spot lesions

Grade 2: severe white spot lesions

Grade 3: white spot lesions with cavitation

2. Determination of fluorescence, size and lesion depth and the bacterial activity of white spot lesions by means of quantitative light-induced fluorescence measurement (QLF) at baseline, after 6 months, and after 12 months of orthodontic treatment.

Secondary outcome measures

Determination of fluorescence, size and lesion depth and the bacterial activity of white spot lesions by means of quantitative light-induced fluorescence measurement (QLF) after active orthodontic treatment and 6 month later.

Overall study start date

01/11/2014

Completion date

01/10/2016

Eligibility

Key inclusion criteria

1. Orthodontic indication for a vestibular multi-bracket appliance or
2. The need and desire to receive Invisalign therapy
3. Persons aged 13-19 years (teenagers) with fully erupted and visible teeth before treatment
4. No white spot lesions at baseline/no caries
5. Good oral hygiene
6. Written informed consent

Participant type(s)

Patient

Age group

Child

Lower age limit

13 Years

Upper age limit

19 Years

Sex

Both

Target number of participants

60

Key exclusion criteria

1. Patients with cheilognathopalatoschisis or other syndromes associated with craniofacial abnormalities
2. Patients with enamel abnormalities

Date of first enrolment

01/11/2014

Date of final enrolment

01/10/2016

Locations

Countries of recruitment

Germany

Study participating centre**Head of the Department of Orthodontics**

Freiburg, Germany

Germany

79106

Sponsor information

Organisation

Align Technology, Inc. (USA)

Sponsor details

2560 Orchard Parkway, San Jose CA 95131

San Jose

United States of America

95131

Sponsor type

Industry

ROR

<https://ror.org/005sbaa41>

Funder(s)

Funder type

Industry

Funder Name

International Align Research Award of the Align Technology, North America (USA)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration