

Effects of structured nutrition education on maternal perceptions, and exclusive breastfeeding duration in Kiandutu health centre, Thika - Kenya

Submission date 02/02/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 21/04/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 06/06/2023	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Breastfeeding has been shown to be important for the health, nutrition and survival of children. This is especially the case in developing countries, where there are high rates of infectious diseases and a lack of good nutrition. Despite this, exclusive breastfeeding rates in Kenya are low, particularly in city slums. This study is looking at a personalized nutrition education program that encourages exclusive breast feeding (EBF) for the first six months and is tailored to improve mothers' breastfeeding self-efficacy. The aim of this study is to look at the effectiveness of this program at improving EBF rates and breast feeding self-efficiency.

Who can participate?

Healthy pregnant women.

What does the study involve?

Participating villages are randomly allocated to one of two groups. Those in the first group take part in the education program. This involves listening to a talk at the health centre about the importance of a proper diet during pregnancy and while breast feeding, and information about the value of breastfeeding to children's health. Those in the second group continue as normal for the duration of the study. Participants in both groups are followed up twice during pregnancy and once a month after they have given birth for six months to find out how many women are exclusively breastfeeding their babies and their views about breastfeeding.

What are the possible benefits and risks of participating?

Participants who take part in the education program benefit from learning about the importance of a good diet and breastfeeding, which could encourage them to eat more healthily and to breastfeed their baby. There are no direct risks involved with participating.

Where is the study run from?

1. Kianhutu Health Centre (Kenya)
2. Makongeni Health Centre (Kenya)

When is the study starting and how long is it expected to run for?

February 2012 to October 2014

Who is funding the study?

National Commission of Science Technology and Innovation (Kenya)

Who is the main contact?

Ms Dorothy Mituki

Contact information

Type(s)

Public

Contact name

Ms Dorothy Mituki

Contact details

Egerton University
Department of Human Nutrition
P.O BOX 536
Egerton
Nakuru
Kenya
20115

Additional identifiers

Protocol serial number

EU/DVRE/028

Study information

Scientific Title

Effects of structured nutrition education on maternal perceptions, and exclusive breastfeeding duration in Kiandutu health centre, Thika - Kenya: A cluster randomized controlled study

Study objectives

Null hypotheses:

1. There is no significant difference in socio-economic, household food security and post-partum factors between mothers in the structured nutrition education group and the comparison group
2. There is no significant difference in BSE and breastfeeding perceptions for mothers in the two groups at baseline
3. There is no significant difference in BSE and breastfeeding perceptions for mothers in the two groups before and after the intervention
4. There is no significant difference in BSE and breastfeeding perceptions for mothers in the

intervention and those in the comparison groups after the intervention

5. There is no significant relationship between socio-economic, household food security post-partum factors, BSE, breastfeeding perceptions and EBF duration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Egerton University Ethics Committee, 07/06/2013, ref: EU/DVRE/028

Study design

Cluster randomised controlled trial

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Breastfeeding

Interventions

The participating villages are randomized to one of two groups in a 1 :1 ratio using the Microsoft excel function.

Intervention group: Participants receive personalized nutrition education at the health facility. This involves a talk on the importance of proper diet during pregnancy and lactation, including dietary diversity, the value of exclusive breastfeeding and the fact that all women can practice EBF with a positive mind set. It will also involve giving them information on how to deal with issues relating to breastfeeding. The content is developed from the Kenyan maternal infant and young child guidelines (which is developed from the UNICEF/WHO maternal infant and young child guidelines). Participants are followed up in their homes by Community Health Workers (CHWs) for 2 times during pregnancy at ≤ 28 and 37 weeks gestation and every month after delivery up to six months post-partum.

Comparison group: Participants receive standard care for the duration of the study. This involves group nutrition education and general health issues done by the health care workers at the facility

Follow up for all participants involves taking their 24 hour recall and weight data at 1, 6, 10 and 14 weeks and 6 months at the study facility.

Intervention Type

Behavioural

Primary outcome(s)

Exclusive breastfeeding rate is measured using the 24 hour recall at 1, 6, 10 and 14 weeks and 6 months.

Key secondary outcome(s))

1. Breastfeeding self-efficacy is measured using the breast feeding self efficacy scale (short form) (BSES-SF) at baseline (34 weeks) at mid (37 weeks) and at end of intervention (6 months)
2. Perceptions towards exclusive breastfeeding is measured using the Deborah McCarter-Spaulding perceptions scale at baseline (34 weeks) at mid (37 weeks) and at end of intervention (6 months)

Completion date

05/10/2014

Eligibility

Key inclusion criteria

1. Women who are less than six months gestation
2. No history of chronic disorders such as hypertension, diabetes, HIV and tuberculosis
3. Aged 18 years and over

Participant type(s)

Other

Healthy volunteers allowed

No

Age group

Other

Lower age limit

18 years

Sex

Female

Total final enrolment

526

Key exclusion criteria

1. Preterm delivery and Infants delivered less than 37 weeks gestation
2. Multiple births, only mothers with singletons continued in the study
3. Low birth weight (below 2500gms)
4. Those who give birth to children born with congenital disorders

Date of first enrolment

03/09/2013

Date of final enrolment

02/10/2014

Locations

Countries of recruitment

Kenya

Study participating centre
Kianhutu Health Centre
Thika-Garissa Highway
Off Broadways Primary School
P.O BOX 3,304
Madaraka Thika
Kenya
01002

Study participating centre
Makongeni Health Centre
Thika-Garissa Highway
P.O BOX 1,747
Madaraka Thika
Kenya
01002

Sponsor information

Organisation
National Commission of Science Technology and Innovation

ROR
<https://ror.org/03wzc0b85>

Funder(s)

Funder type
Government

Funder Name
National Commission of Science Technology and Innovation

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		28/04/2020	06/06/2023	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes