

# Randomised trial of tension-free vaginal tape and transobturator tape as treatment for urinary stress incontinence in women

**Submission date**  
16/09/2005

**Recruitment status**  
No longer recruiting

☐ Prospectively registered

☐ Protocol

**Registration date**  
11/11/2005

**Overall study status**  
Completed

☐ Statistical analysis plan

☒ Results

**Last Edited**  
13/09/2013

**Condition category**  
Urological and Genital Diseases

☐ Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Mr Christopher Mayne

### Contact details

Directorate of Women's, Perinatal & Sexual Health  
Leicester General Hospital  
Gwendolen Road  
Leicester  
United Kingdom  
LE5 4PW  
[christopher.mayne@uhl-tr.nhs.uk](mailto:christopher.mayne@uhl-tr.nhs.uk)

## Additional identifiers

### Protocol serial number

9452

## Study information

### Scientific Title

**Study objectives**

To compare the cure rate and complication rate of the tension-free vaginal tape (TVT) and tension free vaginal tape obturator (TVT-O) procedure (TVT & TVT-O, Gynecare, Somerville, New Jersey)

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Randomised controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Urodynamic stress incontinence

**Interventions**

TVT versus TVT-O

**Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome(s)**

1. Objective cure rate (24 hour pad test)
2. Subjective cure rate (Leakage on 3 day voiding diary)

**Key secondary outcome(s)**

1. Change in quality of life (Kings Health Questionnaire [KHQ])
2. Change in symptom severity (Bristol Female Lower Urinary Tract Symptom Questionnaire [BFLUTS], International Consultation on Incontinence questionnaire [ICIQ])
3. Pre and postoperative complications
4. Pain at 1 hour and 1 week (Visual Analogue Scale [VAS])

**Completion date**

01/10/2006

**Eligibility****Key inclusion criteria**

1. Urodynamic stress incontinence
2. Primary continence procedure

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

Female

**Key exclusion criteria**

1. Detrusor overactivity
2. Voiding Dysfunction
3. Prolapse  $\geq$  pelvic organ prolapse quantification system (POP-Q) stage 2

**Date of first enrolment**

01/10/2004

**Date of final enrolment**

01/10/2006

**Locations****Countries of recruitment**

United Kingdom

England

**Study participating centre**

Directorate of Women's, Perinatal & Sexual Health

Leicester

United Kingdom

LE5 4PW

**Sponsor information****Organisation**

University Hospitals of Leicester NHS Trust (UK)

ROR

<https://ror.org/02fha3693>

## Funder(s)

**Funder type**

Government

**Funder Name**

University Hospitals of Leicester NHS Trust (UK)

## Results and Publications

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/04/2011		Yes	No