

To test whether the anthroposophical drug Stibium D6 has beneficial effect on blood clotting in patients undergoing transurethral resection of the prostate

Submission date 02/07/2012	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 30/07/2012	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 30/07/2012	Condition category Haematological Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

We are carrying out a study of 136 patients who are scheduled for transurethral resection of the prostate (surgical procedure that involves cutting away a section of the prostate gland). During this operation, there is a high risk for bleeding. Therefore, we will test the drug Stibium D6 in order to evaluate its benefit of reduction of bleeding complications. We also test the blood clotting time in 20min intervals during the operation.

Who can participate?

We aim to recruit 136 men, age > 18 years who are scheduled for a transurethral resection of the prostate.

What does the study involve?

Patients will be randomly allocated to receive Stibium D6 intravenous (i.v.) or placebo (dummy). During the operation (under anaesthesia), four blood samples are taken at the beginning of the operation as well as after every 20min. Another blood sample will be taken 1 and 2 days after surgery, respectively. A control examination in our outpatient clinics will be 2 weeks after the operation.

What are the possible benefits and risks of participating?

There could be a direct benefit to those taking part and getting the real drug regarding bleeding complications. If the drug has a beneficial effect on blood clotting, there should be benefits to future patients undergoing transurethral resection of the prostate. Eventually, the drug could be used in surgical interventions other than transurethral resection of the prostate..

The main risk of the intervention is the extra samples of blood that has to be taken. However, the amount of blood is very small. There is no risk for iron deficiency and the related anaemia because of the blood samples. So far, there is no known risk of administering Stibium D6.

Where is the study run from?

The study has been set up by the Urology Department of the University of Bern.

When is study starting and how long is it expected to run for?

It is anticipated that recruitment will start July 2012. Participants will be enrolled on the study for a period of two years.

Who is funding the study?

University Clinic of Urology (Urologische Universitaetsklinik), Switzerland

Who is the main contact?

Dr Beat Roth

urology.berne@insel.ch

Contact information

Type(s)

Scientific

Contact name

Dr Beat Roth

Contact details

Urologische Universitaetsklinik

Inselspital

Bern

Switzerland

3010

Additional identifiers

Protocol serial number

DR2046

Study information

Scientific Title

Double blinded, randomized, placebo-controlled trial to evaluate the efficacy of Stibium D6 on clotting in patients undergoing transurethral resection of the prostate (TURP)

Study objectives

The anthroposophical drug Stibium D6 has beneficial effect on blood clotting in patients undergoing TURP

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethical Committee of the Canton Bern, Switzerland, 03/2012, ref: 235/10

Study design

Randomized double-blinded placebo-controlled single center trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Clotting disorder / intraoperative bleeding / transurethral resection of the prostate

Interventions

Transurethral resection of the prostate in all patients.

50% of patients will receive placebo during this intervention, 50% will receive Stibium D6 intravenous (i.v.).

Stibium D6 -dose: 10ml of 0.000001%, Stibium D6 in 250ml 0.9% NaCl

Total duration of intervention: approximately 1 hour

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Stibium D6

Primary outcome(s)

1. Complications (especially bleeding complications), bleeding complications are measured during the first 14 postoperative days (final evaluation during the final outpatient visit 14 days following surgery)
2. Blood clotting time measured at the beginning of the operation, after 20, 40 and 60 minutes of operation as well as on the 1st and 2nd postoperative day.

Key secondary outcome(s)

1. Intraoperative bleeding score measured during operation
2. Readmissions to hospital evaluated within the first 30 postoperative days
3. Duration of TURP
4. Blood glucose levels measured during operation
5. Duration of catheter in place measured within the first 2 postoperative days (during hospitalisation).
6. Time of postoperative bladder flushing required measured within the first 2 postoperative days (during hospitalisation)

Completion date

30/06/2014

Eligibility

Key inclusion criteria

1. Male
2. >18 years
3. Written informed consent
4. Scheduled for transurethral resection of the prostate

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Peripheral artery occlusive disease
2. Coronary heart disease
3. Anticoagulation therapy (e.g. coumarins)
4. History of stroke
5. Clotting disorder
6. Allergy / intolerance to Stibium D6

Date of first enrolment

01/07/2012

Date of final enrolment

30/06/2014

Locations

Countries of recruitment

Switzerland

Study participating centre

Urologische Universitätsklinik

Bern

Switzerland

3010

Sponsor information

Organisation

University Clinic of Urology (Urologische Universitaetsklinik) (Switzerland)

ROR

<https://ror.org/01q9sj412>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

University Clinic of Urology (Urologische Universitaetsklinik) (Switzerland)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration