

Cardiovascular remodeling in small-for-gestational-age fetuses

Submission date 16/06/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 02/08/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 08/08/2017	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Small for gestational age (SGA) is commonly defined by an ultrasound based estimated weight below the 10th percentile. These fetuses were long considered to be small babies with good post birth outcomes. Clinical studies over the last decade have demonstrated that on average SGA have poorer perinatal results, less than ideal neurodevelopment and higher postnatal cardiovascular risks compared with adequate for gestational age (AGA) newborns (newborns born with normal weight). This data suggests that SGA fetuses are in reality forms of late-onset fetal growth restriction (FGR) where placental insufficiency is not reflected in the Doppler. Consequently, there is a need to develop additional markers of poor perinatal outcome allowing identify late-onset FGR among SGA fetuses. Newer findings suggest that FGR induces cardiac changes that could explain the increased predisposition to cardiovascular disease in adult life even in the FGR subgroups with normal fetal Doppler. The aim of this study is to develop a way to predict mean blood pressure of babies at one year of life using parameters of cardiac function and to correlate birth outcome and other diagnostic antenatal parameters.

Who can participate?

Mothers who are at 32 weeks gestational age who are suspected to have a small baby.

What does the study involve?

Participating mothers are followed each for two to three weeks until delivery using ultrasounds, echocardiography, electrocardiography and cardiotocography. During birth, participants receive the normal care. Blood from the umbilical cord is collected and the birth outcomes are recorded. Babies are followed to measure blood pressure and to monitor their heart (using echocardiography and electrocardiography) at 12 months of age.

What are the possible benefits and risks of participating?

Participants may benefit from increased surveillance during pregnancy and their babies first year of life. There are no risks with participating.

Where is the study run from?

Technical University of Munich Department of Obstetrics and Perinatal Medicine, Klinikum rechts der Isar (Germany)

When is the study starting and how long is it expected to run for?
April 2014 to December 2019

Who is funding the study?
Else Kröner-Fresenius Stiftung, EKFS (Germany)

Who is the main contact?
Dr Silvia M. Lobmaier
silvia.lobmaier@tum.de

Contact information

Type(s)
Scientific

Contact name
Dr Silvia M. Lobmaier

ORCID ID
<https://orcid.org/0000-0002-5696-6686>

Contact details
Department of Obstetrics and Gynecology, Obstetrics and perinatal medicine) (Frauenklinik und Poliklinik, Sektion Geburtshilfe und Perinatalmedizin
Technical University of Munich
Ismaninger Str. 22
München
Germany
81675
+49 894 1405417
silvia.lobmaier@tum.de

Additional identifiers

Protocol serial number
CU_001_300

Study information

Scientific Title
Fetal programming: Evaluation of cardiovascular remodeling in late-onset small-for-gestational-age fetuses and prediction of adverse birth and intermediate-term cardiovascular outcome

Acronym
CURIOSA

Study objectives
Study aims:
1. To develop and evaluate a prognostic model for mean RR at one year of age using parameters of cardiac function (average acceleration capacity, TAPSE, right sphericity index, Strain-rate,

aortic distensibility, isovolumetric relaxation time)

2. Evaluate the correlation between the above mentioned diagnostic antenatal parameters and other secondary short- and intermediate-term adverse outcomes; adverse birth outcome [pH < 7.15, operative delivery for fetal distress (fetal scalp blood analysis pH < 7.20 or abnormal fetal heart rate tracing)], abnormal child growth pattern and abnormal metabolic profile. Abnormal cardiovascular outcome at birth and at 12 months of age (aortic intima media thickness > 75th centile and/or RR > 95th centile)

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of the Faculty of Medicine of the Technical University of Munich (Ethikkommission der Fakultät für Medizin der Technischen Universität München), 24/07/2014

Study design

Prospective longitudinal observational study

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Fetal programming, cardiovascular remodeling, late-onset intrauterine growth restriction

Interventions

This study is a prospective longitudinal observational study that includes participants consecutively. All cases are seen at the high risk SGA outpatient ward. The source population is recruited from the population under treatment with term fetuses (Gestational age ≥ 32 weeks at inclusion and born after 37 weeks gestation) suspected of growth restriction defined as an estimated fetal weight or fetal abdominal circumference, measured below the 10th population or customized percentile or a change of abdominal circumference < 5mm in 2 weeks (=reduced growth velocity according to Green Top Guidelines).

Diagnostic tools

During pregnancy: Pregnancies complicated with suspected growth restriction are followed each two to three weeks until delivery in a routine fashion including fetal biometry, advanced ultrasound investigation (including fetal functional echocardiography: Aortic intima media thickness, aortic distensibility, right/Left sphericity index, right/Left cardiac output, tricuspid annular plane systolic excursion (TAPSE), right E/A ratio, tight E', left myocardial performance index, including IRT measurement, strain/strain rate), feto-maternal Doppler, measurement of amniotic fluid and placenta, and registration of fetal heart rate with fetal ECG (AN 24 MONICA device) and simultaneously with conventional computerised CTG for 40 minutes.

Obstetric ultrasound examinations are performed using Voluson E8 or Samsung HS 70A whereas functional echocardiographic parameters are measured using Samsung HS 70 A. All functional echocardiographic parameters are calculated off-line within four weeks after evaluation

according to a standardised protocol based on a trace method with the assistance of commercially available software (TOMTEC ImageArena software). AGA group will be scanned to get 15 controls for each gestational week from 32 to 41 weeks of gestation.

During birth: Participants receive the standard evaluation during labour as do other pregnancies complicated by FGR. Fetal cord blood is taken after cord clamping for metabolic analysis. Placenta is sent to examination by a stated placental pathologist. Birth outcomes are recorded.

Participating infants are followed up for a postnatal cardiovascular evaluation using a blood pressure measurement and echocardiography as well as electrocardiography at 12 months of age.

For evaluation of normal pattern of newer fetal echocardiography parameter and metabolic profile, a group of gestational-age matched (AGA) controls will be examined.

Intervention Type

Mixed

Primary outcome(s)

Prediction of mean blood pressure (BP) (in mmHg) at one year of age is measured using the average acceleration capacity (bpm), tricuspid annular plane systolic excursion (mm), right sphericity index, Strain-rate, aortic distensibility, isovolumetric relaxation time (ms)) during pregnancy.

Key secondary outcome(s)

1. Adverse birth outcomes are measured using
 - 1.1. Mortality defined as intra-uterine death or death before hospital discharge
 - 1.2. Acidosis defined as arterial cord blood pH<7.15
 - 1.3. Apgar at 5 minutes <7
 - 1.4. Admission to NICU
 - 1.5. Operative delivery for fetal distress (fetal scalp blood analysis pH < 7.20 or abnormal fetal heart rate tracing
2. Abnormal child growth pattern are measured using patient records (as defined below the 10th percentile of normal population) during first year of life
3. Abnormal metabolic profile (umbilical cord blood) at birth in comparison to the control group
4. Abnormal cardiovascular outcomes are measured using aortic intima media thickness >75th centile and/or BP > 95th centile at birth and at 12 months of age
5. Prediction of mean blood pressure (BP) at one year of age (mm Hg) using the following measurements: metabolic profile, fetal functional echocardiographic parameters others than the ones used for primary outcome
6. Correlation of mean blood pressure (BP) at one year of age (mm Hg) and autonomic nervous system [average acceleration and deceleration capacity (bpm)] and antenatal/ fetal functional echocardiographic parameters
7. Correlation of antepartal cardiac function parameters and those at 1 year of life

Completion date

31/12/2019

Eligibility

Key inclusion criteria

1. Pregnant women aged 18 and older
2. Fetuses with a gestational age ≥ 32 weeks
3. Suspected of growth restriction defined as an estimated fetal weight or fetal abdominal circumference measured below the 10th population or customized percentile or a change of abdominal circumference $<5\text{mm}$ in 2 weeks measured by ultrasound

Healthy volunteers allowed

No

Age group

All

Sex

All

Key exclusion criteria

1. Born <37 weeks gestational age
2. Known chromosomal and/or structural anomaly
3. Multiple gestation
4. Mothers with cardiac heart disease
5. Mothers' severe systemic disease
6. Antihypertensive medication
7. Antenatal intra-uterine infection

Date of first enrolment

01/09/2016

Date of final enrolment

30/09/2019

Locations**Countries of recruitment**

Germany

Study participating centre

Frauenklinik und Poliklinik, Klinikum rechts der Isar of the Technical University of Munich

Ismaninger Str. 22

München

Germany

81675

Sponsor information**Organisation**

Else Kröner-Fresenius Foundation (Else Kröner-Fresenius-Stiftung)

ROR

<https://ror.org/03zcxha54>

Funder(s)

Funder type

Charity

Funder Name

Else Kröner-Fresenius-Stiftung

Alternative Name(s)

Else Kroner-Fresenius Foundation, Else Kroener-Fresenius-Stiftung, Else Kröner Fresenius-Stiftung, EKFSStiftung, StiftungEKFS, EKFS

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

Germany

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from PD Dr. Silvia Lobmaier, Section of Obstetrics and Perinatal Medicine of the Department of Gynecology of Klinikum rechts der Isar, Technical University of Munich, Germany

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes