# Radioimmunotherapy with 90Y Ibritumumab Tiuxetan in B-cell CD20+ indolent non-Hodgkin Lymphoma

Submission date	Recruitment status	Prospectively registered
17/03/2013	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
10/04/2013	Completed	[X] Results
Last Edited	Condition category	Individual participant data
08/01/2015	Cancer	

## Plain English summary of protocol

Background and study aims

Treatment with 90YIbritumomab/tiuxetan (90Y-IT) in CD20+ for non Hodgkin lymphoma (NHL) patients is safe and effective. We have reviewed our hospital database to analyse the outcomes of CD20+ NHL patients over 65 years old treated with 90Y-IT

Who can participate?

Patients diagnosed with CD20+ indolent NHL.

What does the study involve?

To analyze the results of therapy with 90Y-Ibritumomab/tiuxetan in clinical practice prescription.

What are the possible benefits and risks of participating?

90Y-IT has demonstrated high efficacy and safety in the therapy of CD20+ indolent NHL patients. We want to analyse these aspects in patients treated over 65 years old and to evaluate safety, survival and comorbidities in the long term.

Where is the study run from?

The Miguel Servet University Hospital, Zaragoza (Spain).

When is the study starting and how long is it expected to run for?

The study started in September 2005 when the first patient received therapy with 90Y-IT in the Miguel Servet University Hospital.

Who is funding the study?

The Foundation for the Study of Hematology in Aragon (FEHHA)

Who is the main contact? Dr Pilar Giraldo fehha@fehha.org

## Contact information

## Type(s)

Scientific

#### Contact name

Dr Pilar Giraldo

#### Contact details

Haematology Department Research Traslational Unit Miguel Servet University Hospital Zaragoza Spain 50009

## Additional identifiers

### Protocol serial number

**RIT-002** 

## Study information

#### Scientific Title

Radioimmunotherapy with 90Y Ibritumumab Tiuxetan in B-cell CD20+ indolent non-Hodgkin Lymphoma: A retrospective study

#### Acronym

RIT

## **Study objectives**

Radioimmunotherapy (RIT) emerged as an option after Mo-Ab success, combining the selectivity of the anti CD-20 immunotherapy with an attached radioisotope, making possible the delivery of radiation exactly within the tumor burden. The previous phase III randomized study that compared 90Y-IT with Rituximab immunotherapy in relapsed or refractory low-grade follicular or transformed CD20+ NHL showed significant ORR for the 90Y-IT group vs the Rituximab group. In addition the results of the phase III randomized study of 90Y-IT as front-line consolidation, the First-line Indolent Trial (FIT) published in 2008, showed that 90Y-IT consolidation significantly prolonged median Progression Free Survival (PFS) after a median of 3.5 years of observation, and converted 77% of Partial Response (PR) after induction into CR, resulting in a CR rate of 87% with low associated toxicity.

We incorporated RIT with 90Y Ibritumumab Tiuxetan in B-cell CD20+ indolent non-Hodgkin Lymphoma since 2005 according an a clinical protocol conducted by a multidisciplinary team formed by clinical hematologists, nuclear medicine physicians, radiopharmacy physicians and nurses in the Miguel Servet University Hospital at Saragossa, Spain. In 2007 the therapeutic inclusion was extended as consolidation after first-line chemotherapy in patients with complete or partial response confirmed by PET/CT. All patients had been treated and follow-up with the same criteria and now we want to analyze retrospectively the results after long term follow-up.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Ethics Committee Comite de estudios de investigación clínica de Aragón (CEICA),

## Study design

Interventional non-randomised retrospective study

## Primary study design

Interventional

## Study type(s)

**Treatment** 

## Health condition(s) or problem(s) studied

B-cell CD20+ indolent non-Hodgkin Lymphoma

#### Interventions

A retrospective study based in a clinical protocol conducted by a multidisciplinary team comprising clinical hematologists, nuclear medicine physicians, radiopharmacists, and nurses at MSUH

Therapy with 90-YIbritumomab/tiuxetan in relapsed and refractory indolent B-cell NHL and in consolidation after first-line immuno- or chemoimmunotherapy.

The study is open in one arm and all patients received an intravenously infusion of Rituximab on day 1 at dose of 250 mg/m2 and a second dose of Rituximab 250 mg/m2 on day 7 followed by weight-based dose of 90Y-IT administered as slow intravenous push over 10 minutes within the next 4 hours of Rituximab infusion.

## Intervention Type

Drug

#### Phase

Phase IV

## Drug/device/biological/vaccine name(s)

90Y Ibritumumab Tiuxetan

## Primary outcome(s)

Response assessment was made 12 weeks after treatment

PET/CT scan was performed in all cases and response criteria used were the same as the International Working Group (IWG). and classified as Complete Response (CR), Partial Response (PR) and no response (NR).

## Key secondary outcome(s))

- 1. Time to Progression (TTP)
- 2. Overall Response (OR) and OS in all patients.
- 3. We also registered side effects, with special emphasis in myelotoxicity and emerging second neoplasms.

TTP and OS were calculated from the date of 90Y-IT therapy until disease progression or death. All eligible patients were accepted by the clinical committee and included into the analysis.

### Completion date

29/02/2012

## **Eligibility**

## Key inclusion criteria

- 1. Adults patients with an excisional biopsy-confirmed diagnosis of CD20+ F-NHL grade 1, 2, or 3a according to the revised WHO classification system
- 2. ECOG performance status  $\leq 2$
- 3. Absolute neutrophil count (ANC) ≥ 1,500/µL
- 4. Absolute platelet count (APC) ≥ 100,000/µL
- 5. Bone marrow total lymphocytes ≤ 25% by morphological counting
- 6. Serum bilirubin  $\leq 2.0 \text{mg/dL}$  and serum creatinine  $\leq 2.0 \text{mg/dL}$
- 7. All patients were requested to sign an informed consent form

## Participant type(s)

Patient

## Healthy volunteers allowed

No

### Age group

Adult

#### Sex

All

### Kev exclusion criteria

- 1. Presence of any other malignancy or history of prior malignancy within 5 years of study entry. This does not apply to-non melanoma skin tumors and in situ cervical cancer.
- 2. Prior radioimmunotherapy
- 3. Presence of primary central nervous system (CNS) lymphoma at first diagnosis
- 4. Known seropositivity for hepatitis C virus (HCV) or hepatitis B surface antigen (HbsAg).
- 5. Known history of HIV infection
- 6. Abnormal liver function: total bilirubin > 2 × ULN unless secondary to Gilbert disease.
- 7. Abnormal renal function: serum creatinine > 2.0 × ULN
- 8. Ongoing toxic effects of chemotherapy > grade 2 and expected to interfere with Zevalin® treatment.

#### Date of first enrolment

01/09/2005

# Date of final enrolment 29/02/2012

## Locations

Countries of recruitment

Spain

Study participating centre Haematology Department Zaragoza Spain 50009

# Sponsor information

## Organisation

Foundation for the Study of hematology in Aragon [Fundación para el Estudio de la Hematologia y Hemoterapia en Aragón](FEHHA)

# Funder(s)

## Funder type

Research organisation

#### **Funder Name**

Foundation for the Study of hematology in Aragon [Fundación para el Estudio de la Hematologia y Hemoterapia en Aragón](FEHHA) (Spain)

## **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type

**Details** results

Date created Date added Peer reviewed? Patient-facing?

Results article 01/12/2014 Yes No

Participant information sheet Participant information sheet 11/11/2025 No Yes