

# Comparing pain relief methods for keyhole bowel surgery: A study on two types of nerve blocks

<b>Submission date</b> 18/11/2024	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 25/11/2024	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 23/09/2025	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

This study aims to compare two pain relief methods used after keyhole bowel surgery. The goal is to find the most effective way to reduce pain and minimize the need for morphine, which can have side effects. The two methods being compared are the quadratus lumborum (QL) block and the transversus abdominis plane (TAP) block.

### Who can participate?

Patients who are undergoing laparoscopic (keyhole) colorectal surgery can participate in this study.

### What does the study involve?

Participants will be randomly assigned to one of two groups: one group will receive the QL block, and the other will receive the TAP block. Both groups will receive general anesthesia during surgery. After the surgery, participants will receive one of the two pain relief methods. All participants will also receive standard pain relief medications, including paracetamol, nefopam, and patient-controlled analgesia (PCA).

### What are the possible benefits and risks of participating?

Participants may benefit from effective pain relief from one of the two methods. Both methods have been shown to reduce pain after laparoscopic colorectal surgery. There is a very small risk of complications related to anesthesia, but these are rare and will be closely monitored by medical staff.

### Where is the study run from?

Binh Dan Hospital (Viet Nam)

### When is the study starting and how long is it expected to run for?

July 2024 to October 2025

Who is funding the study?  
University of Medicine and Pharmacy at Ho Chi Minh City Medicine and Pharmacy University (Viet Nam)

Who is the main contact?  
Mr Van Phuoc Toan, dr.toanvan@gmail.com

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Mr Phuoc Toan Van

### ORCID ID

<https://orcid.org/0009-0009-6233-0826>

### Contact details

371 Dien Bien Phu - Ward 4 - District 3  
Ho Chi Minh  
Viet Nam  
700000  
+84 902993591  
dr.toanvan@gmail.com

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### ClinicalTrials.gov (NCT)

Nil known

### Protocol serial number

Nil known

## Study information

### Scientific Title

Comparison of analgesic efficacy between quadratus lumborum block and transversus abdominal plane block in laparoscopic colorectal surgery

### Study objectives

Quadratus lumborum anesthesia reduces morphine consumption 48 hours postoperatively by 50% compared with transverse abdominal plane anesthesia in laparoscopic colorectal surgery

### Ethics approval required

Ethics approval required

**Ethics approval(s)**

approved 22/10/2024, Ethics council in biomedical research, University of Medicine and Pharmacy at Ho Chi Minh city (217 Hong Bang, Cho Lon Ward, District 5, Hochiminh city, 700000, Viet Nam; +84 28 3855 0507; hoidongdaoducdhyd@ump.edu.vn), ref: IRB-VN01002/IRB00010293 /FWA00023448

**Study design**

Randomized controlled trial

**Primary study design**

Interventional

**Study type(s)**

Efficacy

**Health condition(s) or problem(s) studied**

Analgesic after laparoscopic colorectal surgery

**Interventions**

All patients recruited into the study will be randomized into 2 groups: GL group (quadra lumbar muscle block group) and TAP group (transversus abdominis plane block group) by creating a block random number table in Excel software 2016 (copyrighted by the Department of Public Health - Ho Chi Minh City University of Medicine and Pharmacy). Choosing the RAND function with the sort A->Z command will give you similar random numbers corresponds to the serial number of the patients in the research sample. Choose an odd number QL group and even numbers for TAP group.

The patient is induced, intubated and maintained according to the protocol hospital. At the end of surgery: After the last stitch of skin is sutured, the patient is fine perform QL or TAP according to initial randomization.

Both methods are performed under ultrasound guidance. The anesthetic used is ropivacaine 0.25% 20ml for each side.

Post-operative analgesia includes: patient-controlled analgesia (PCA) with morphine, paracetamol 1gr per 8h, nefopam 20mg per 8h.

**Intervention Type**

Procedure/Surgery

**Primary outcome(s)**

Total dose of morphine-PCA 48 hours after surgery measured using patient records

**Key secondary outcome(s)**

Measured using patient records:

1. Total dose of morphine-PCA at postoperative time points (1 hour, 2 hours, 6 hours, 12 hours, 24 hours)
2. VAS pain scores at rest and movement at postoperative times (1 hour, 2 hours, 6 hours, 12 hours, 24 hours, 48 hours)
3. Postoperative nausea and vomiting (PONV)
4. Opioid sedation (POSS)
5. Abdominal skin sensory area after 1 hour of blockade.

**Completion date**

30/10/2025

## Eligibility

### Key inclusion criteria

Current inclusion criteria as of 23/09/2025:

1. The patient is indicated for elective laparoscopic colorectal surgery
2. Patients are 18 to 75 years old
3. Patients' health status is assessed according to the Society of Anesthesiology and Resuscitation United States (ASA) at level I-III

---

Previous inclusion criteria:

1. The patient is indicated for elective laparoscopic colorectal surgery
2. Patients are 18 to 70 years old
3. Patients' health status is assessed according to the Society of Anesthesiology and Resuscitation United States (ASA) at level I-III

### Participant type(s)

Resident

### Healthy volunteers allowed

No

### Age group

Adult

### Lower age limit

18 years

### Upper age limit

75 years

### Sex

All

### Total final enrolment

60

### Key exclusion criteria

1. Patients addicted to opioid drugs
2. Do not understand the instructions for using the PCA device
3. History of chronic pain, anesthetic allergy
4. Obese patients have BMI over 30 kg/m<sup>2</sup>
5. The patient converted from laparoscopic surgery to open surgery

### Date of first enrolment

11/11/2024

**Date of final enrolment**

30/06/2025

## **Locations**

**Countries of recruitment**

Viet Nam

**Study participating centre**

**Binh Dan Hospital, Hochiminh city**

371 Dien Bien Phu - Ban Co Ward

Hochiminh city

Viet Nam

700000

## **Sponsor information**

**Organisation**

Binh Dan Hospital

## **Funder(s)**

**Funder type**

University/education

**Funder Name**

University of Medicine and Pharmacy at Ho Chi Minh City Medicine and Pharmacy University

## **Results and Publications**

**Individual participant data (IPD) sharing plan**

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

hoinghibvbd@gmail.com

dr.toanvan@gmail.com

**IPD sharing plan summary**

Stored in non-publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>			23/09/2025	No	Yes
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes