Assessing airway twitchiness in patients with severe asthma on biologics as they reduce their inhaled corticosteriod dose

| Submission date | Recruitment status Recruiting | [X] Prospectively registered | | |
|-------------------|--------------------------------------|---------------------------------|--|--|
| 15/07/2025 | | [X] Protocol | | |
| Registration date | Overall study status | Statistical analysis plan | | |
| 16/07/2025 | Ongoing Condition category | ☐ Results | | |
| Last Edited | | Individual participant data | | |
| 16/07/2025 | Respiratory | [X] Record updated in last year | | |

Plain English summary of protocol

Background and study aims

Asthma is a common lung condition that makes breathing difficult, especially when the airways become sensitive or "twitchy" in response to things like dust or pollen. This twitchiness is linked to a type of inflammation in the lungs called type 2 inflammation. People with this kind of asthma often have higher levels of certain markers in their blood and breath.

Asthma is usually treated with inhalers that contain steroids, but these don't always reach the smallest parts of the lungs. Newer treatments called biologics, which are given by injection, can reach these areas and may help reduce inflammation more effectively.

This study aims to find out whether two commonly used biologic medications, dupilumab and tezepelumab, can reduce airway twitchiness enough that patients can safely lower their dose of steroid inhalers without losing control of their asthma.

Who can participate?

The study is for adults with severe asthma who are already being treated with either dupilumab or tezepelumab.

What does the study involve?

Participants will use a standard steroid inhaler in a specific way called maintenance and reliever therapy. They will attend three study visits over six months, including a screening visit. They will provide blood samples to measure inflammation markers and take breathing tests to check lung function, including a special test to measure how twitchy the airways are.

What are the possible benefits and risks of participating?

Benefits may include better understanding and management of asthma, and possibly reducing the steroid dose safely. Risks include side effects from reducing steroid use, such as worsening asthma symptoms. There are also known side effects from long-term use of high-dose steroid inhalers, which this study hopes to help reduce. Blood tests and breathing tests may cause minor discomfort.

Where is the study run from?

The study is being run at the Scottish Centre of Respiratory Medicine, based at the University of Dundee's Ninewells campus (UK)

When is the study starting and how long is it expected to run for? May 2024 to August 2026

Who is funding the study? Investigator initiated and funded

Who is the main contact? Professor Brian Lipworth, b.j.lipworth@dundee.ac.uk

Contact information

Type(s)

Principal Investigator

Contact name

Prof Brian Lipworth

ORCID ID

https://orcid.org/0000-0002-8140-2014

Contact details

Scottish Centre for Respiratory Research Mailbox 2
Molecular & Clinical Medicine School of Medicine
University of Dundee Ninewells Hospital
Dundee
United Kingdom
DD1 9SY
+44 1382383188
b.j.lipworth@dundee.ac.uk

Type(s)

Public, Scientific

Contact name

Dr Robert Greig

ORCID ID

https://orcid.org/0000-0001-7668-286X

Contact details

Scottish Centre for Respiratory Research Mailbox 2 Molecular & Clinical Medicine School of Medicine University of Dundee Ninewells Hospital Dundee United Kingdom DD1 9SY

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

346249

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

2-071-24

Study information

Scientific Title

Tailoring Inhaled Corticosteroids in patients with Severe Asthma taking Biologics

Acronym

TICSAB

Study objectives

- 1. To characterize the effect of tapering inhaled corticosteroids on mannitol airway hyperresponsiveness in patients with severe asthma taking dupilumab or Tezepelumab
- 2. Assess the effects of taking dupilumab or tezepelumab on small airways dysfunction using airways oscillometry.
- 3. To characterise symptom control and quality of life scores as participants adjust their inhaled corticosteroid dose.
- 4. To assess type 2 inflammation in as participants adjust their inhaled corticosteroid dose.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 08/07/2025, East of Scotland Research Ethics Service (EoSRES) (Tayside Medical Science Centre, Residency Block Level 3, George Pirie Way, Ninewells Hospital and Medical School, Dundee, DD1 9SY, United Kingdom; +44 1382660111; tay.eosres@nhs.scot), ref: 25/ES/0043

Study design

Phase IV single arm open labelled

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Home, Hospital

Study type(s)

Quality of life, Treatment

Participant information sheet

See study outputs table

Health condition(s) or problem(s) studied

Severe asthma patient on either dupilumab or tezepelumab

Interventions

Participants will already be established on asthma biologic medications (either dupilumab or tezepelumab). Will will be on maximal maintenance and reliever inhaled therapy (Fostair NEXTHaler 100/6 4 puffs BD) and reduce their MART dose as able during the study (between 2 and 8 puffs daily) over a 6 month period. They will undergo airway oscillometry, spirometry, mannitol challenge, blood tests, FeNO and questionnaires to assess symptom control and quality of life at 3 visits (including a screening visit).

Intervention Type

Other

Primary outcome measure

Mannitol airway hyperresponsiveness measured using change in mannitol PD10 from Visit 1 (post-run-in baseline) to Visit 2 (6 months)

Secondary outcome measures

- 1. Blood eosinophil count is measured using automated haematology analyser at post-run-in baseline and 6 months
- 2. Total serum IgE is measured using immunoassay (e.g. ImmunoCAP) at post-run-in baseline and 6 months
- 3. Fractional exhaled nitric oxide (FeNO) is measured using a FeNO analyser (e.g. NIOX VERO) at post-run-in baseline and 6 months
- 4. Forced expiratory volume in 1 second (FEV1), forced expiratory flow at 25–75% of FVC (FEF25–75), and forced vital capacity (FVC) are measured using spirometry at post-run-in baseline and 6 months
- 5. Airway resistance (R5–R20) and reactance area (AX) are measured using impulse oscillometry at post-run-in baseline and 6 months
- 6. Airway hyperresponsiveness is measured using mannitol challenge test with PD15 as the outcome at 6 months
- 7. Airway reactance area (AX) is measured using impulse oscillometry at post-run-in baseline and 6 months

Overall study start date

01/05/2024

Completion date

01/08/2026

Eligibility

Key inclusion criteria

- 1. Any patient over 18 years of age with severe asthma taking dupilumab or tezepelumab for severe asthma for at least 6 months
- 2. FEV1 ≥50% at baseline

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

100 Years

Sex

Both

Target number of participants

46

Total final enrolment

46

Key exclusion criteria

- 1. Any patients on maintenance oral steroids or required an oral steroid burst in the past 28 days
- 2. Any patient who was switched from another biologic in the past 3 months
- 3. Any other respiratory condition such as moderate to severe bronchiectasis or COPD
- 4. Currently pregnant

Date of first enrolment

01/08/2025

Date of final enrolment

10/01/2026

Locations

Countries of recruitment

Scotland

United Kingdom

Study participating centre Ninewells Hospital and University of Dundee Medical school Ninewells Avenue Dundee United Kingdom DD1 9SY

Sponsor information

Organisation

TASC - University of Dundee/NHS Tayside

Sponsor details

Tayside Medical Science Centre, Ninewells Hospital & Medical School Research & Development Office Residency Block, Level 3, George Pirie Way Dundee Scotland United Kingdom DD1 9SY +44 1382383297 tascgovernance@dundee.ac.uk

Sponsor type

University/education

Website

https://www.dundee.ac.uk/tasc

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Publication and dissemination plan

Planned publication in peer reviewed journal.

Intention to publish date

01/08/2027

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available on request from Prof Brian Lipworth (b.j.lipworth@dundee.ac.uk)

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-------------------------------|-----------|--------------|------------|----------------|-----------------|
| Participant information sheet | version 3 | 30/06/2025 | 16/07/2025 | No | Yes |
| Protocol file | version 2 | 30/06/2025 | 16/07/2025 | No | No |