Effects of transfusing fresh versus standardissue red cells on in-hospital mortality

Submission date	Recruitment status No longer recruiting	Prospectively registered		
04/08/2010		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
11/01/2011		[X] Results		
Last Edited	Condition category	Individual participant data		
24/08/2012	Surgery			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Nancy Heddle

Contact details

McMaster Transfusion Research Program HSC-3H50 1200 Main St West Hamilton Canada L8N 3Z5 +1 905 52591040 ext 22126 heddlen@mcmaster.ca

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Effects of transfusing fresh versus standard-issue red cells on in-hospital mortality: a pilot randomised controlled trial using a pragmatic approach

Acronym

INFORM

Study objectives

Hundreds of thousands of Canadians receive a red cell transfusion each year. Current methods of blood inventory management are to issue the oldest blood first to limit outdating. However, data from experimental and observational studies suggest that red cells stored for longer periods might lead to adverse outcomes, including an increase in mortality. The only way to answer this question is to undertake a randomised controlled trial (RCT). If such a trial showed a statistically significant and clinically important improvement in mortality with freshest available versus standard issue red cells, it is likely to lead to major changes in the management of red cell inventories and storage methods to increase the use of fresher red cells. Such a large pragmatic trial is complex; hence before initiating a large trial it is important to work out logistics and show feasibility. Our proposed pilot study will provide crucial information for the design of a large RCT that will yield reliable and precise estimates of the effect of freshest available versus standard-issue red cells on in-hospital mortality.

Further reading:

Duration of red cell storage before transfusion and in-hospital mortality. Eikelboom JW, Cook RJ, Liu Y, Heddle NM. Am Heart J. 2010 May;159(5):737-743.e1. http://www.ncbi.nlm.nih.gov/pubmed/20435180

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Research Ethics Board at Hamilton Health Sciences approved on the 20th April 2010 (ref: 10-196)

Study design

Single centre pilot pragmatic double blind randomised controlled trial

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Red cell blood transfusion

Interventions

Patients requiring red cell transfusion will be randomised to one of the following conditions:

- 1. Experimental transfusion is freshest available red cells
- 2. Control is standard issue red cells (oldest product in stock)

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

Feasibility of randomising consecutive patients requiring blood transfusion: we anticipate that early in the study the failure rate may be higher due to the learning curve associated with the randomisation process; hence, the feasibility failure rate will be assessed on the data from the final 3 months of the 6 months pilot. This outcome will be monitored weekly with changes being implemented to prevent failures from occurring if possible. This is vital information in determining the feasibility of the larger RCT.

Key secondary outcome(s))

- 1. Impact on inventory and red cell outdating rate
- 2. Contrast in the age of fresh and standard-issue red cells
- 3. Ability to provide timely reports to monitor inventory levels
- 4. Outdating and age overlap

The frequency of in-hospital mortality will also be documented but used only to estimate the sample size for expanding the study if feasibility is demonstrated.

Completion date

01/12/2010

Eligibility

Key inclusion criteria

- 1. Patients who are admitted to Hamilton General Hospital
- 2. Destined to receive a red cell transfusion
- 3. Either sex, no age restrictions

The REB has approved that this study to be done with waived consent as it meets the 5 requirements for waived consent identified by the Tri Council Policy. However, to meet all requirements of this policy, patients will be informed that this study is taking place and will be given a summary of the study.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Other

Sex

Αll

Key exclusion criteria

1. Patients who have a particular requirement for fresh red cells (e.g., sickle cell disease, transfusion dependent thalassemia, fresh cells requested by physician)

- 2. Are to receive pre-planned directed or autologous donations
- 3. Massive transfusion anticipated
- 4. Being transfused as an outpatient

Date of first enrolment

01/01/2010

Date of final enrolment

01/12/2010

Locations

Countries of recruitment

Canada

Study participating centre McMaster Transfusion Research Program Hamilton

Hamilton Canada L8N 3Z5

Sponsor information

Organisation

McMaster University (Canada)

ROR

https://ror.org/02fa3aq29

Funder(s)

Funder type

Research organisation

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - http://www.cihr-irsc.gc.ca (ref: 221072)

Funder Name

Canadian Blood Services (Canada)

Alternative Name(s)

Société canadienne du sang, Canadian Blood Services - Ontario, lifelineontario, CBS

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

Canada

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2012		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes