The pharmacokinetics and potential health effects of champagne wine in human subjects

Submission date	Recruitment status No longer recruiting	Prospectively registered		
09/07/2009		Protocol		
Registration date	Overall study status	Statistical analysis plan		
07/08/2009	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
31/01/2013	Circulatory System			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Jeremy Paul Edward Spencer

Contact details

Department of Food and Nutritional Sciences School of Chemistry, Food Biosciences and Pharmacy The University of Reading PO Box 226, Whiteknights Reading United Kingdom RG6 6AP +44 (0)118 378 8724 j.p.e.spencer@reading.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT) NCT00937313

Protocol serial number

N/A

Study information

Scientific Title

The pharmacokinetics and potential health effects of champagne wine in human subjects: a placebo-controlled randomised cross-over human trial

Study objectives

Determine whether there is a link between champagne wine intake and changes in cardiovascular disease (CVD) risk factors, including blood lipid profile, platelet function and oxidative status. The study will also establish the absorption of polyphenol uptake following consumption of champagne wine.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The University of Reading Research Ethics Committee approved on the 8th May 2007 (ref: 07/16)

Study design

Placebo-controlled randomised cross-over human trial

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Cardiovascular disease (CVD) risk factors

Interventions

Acute consumption of three glasses of champagne wine (375 ml) or the equivalent alcohol with sparkling mineral water as the control (placebo) within a 10-minute period. This amount of champagne contains 4.5 units of alcohol, which is above the legal limit for operating a car or other machinery. Following a washout period of 28 days, volunteers returned to the unit to complete the second arm of the study where the procedure above was repeated.

Subjects were assessed for anthropometric measurements and provided a urine sample prior baseline Laser Doppler Imaging with iontophoresis (LDI) measurements. Subjects were then cannulated and a baseline blood sample was collected. The intervention was then performed. Following a standardised breakfast blood samples were collected at: 15, 30, 45, 60, 120, 180, 240, 300, 360 and 480 minutes post-consumption and pooled urine samples were collected over 3 x 8-hour periods. A standardised breakfast and lunch were also consumed at 15 and 200 minutes post beverage. LDI measurements were carried out at 120, 240, 360 and 480 minutes. Subjects also provided 24-hour and 32-hour blood and urine samples.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

- 1. Assessment of endothelial function by Laser Doppler Imaging with iontophoresis, at baseline and 28 days
- 2. Blood assessment of lipid profile, inflammatory markers, plasma antioxidant and oxidant capacity, liver enzyme and metalloproteinase blood concentrations, measured at baseline, 15, 30, 45, 60, 120, 180, 240, 300, 360 and 480 minutes post-consumption at each intervention point

Key secondary outcome(s))

Bioavailability of phytochemicals and metabolite excretion, pooled urine samples were collected over 3 x 8-hour periods, at 24 hours and 32 hours at each intervention point

Completion date

01/08/2009

Eligibility

Key inclusion criteria

- 1. Healthy male and female subjects
- 2. Aged between 20 and 65 years
- 3. Body mass index (BMI) between 19 and 25 kg/m^2
- 4. Normal concentrations of liver enzymes (aspartate aminotransferase [AST], alanine aminotransferase [ALT], gamma-glutamyl transpeptidase [gamma-GT])
- 5. Normal haemoglobin, hematocrit and leucocyte counts
- 6. Absence of glucose and protein in urine

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

- 1. Diabetes mellitus
- 2. Any form of liver or gastrointestinal disorder
- 3. Low body mass index [BMI] (less than 19 kg/m^2)
- 4. High blood pressure (greater than 150/90 mmHg)
- 5. Anaemia
- 6. Gall bladder problems
- 7. Present illness
- 8. Taking dietary supplements
- 9. Vigorous exercise (greater than 3 x 20 minutes/week)
- 10. Alcohol consumption more than 120 g (women) and 168 g (men) per week
- 11. Pregnant or lactating females

Date of first enrolment

01/09/2008

Date of final enrolment

01/08/2009

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Department of Food and Nutritional Sciences

Reading United Kingdom RG6 6AP

Sponsor information

Organisation

Biotechnology and Biological Sciences Research Council (BBSRC) (UK)

ROR

https://ror.org/00cwqg982

Funder(s)

Funder type

Research council

Funder Name

Biotechnology and Biological Sciences Research Council (BBSRC) (UK) (ref: BB/F008953/1; BB /C518222/1; BB/G005702/1)

Alternative Name(s)

UKRI - Biotechnology And Biological Sciences Research Council, BBSRC UK, Biotechnology and Biological Sciences Research Council (BBSRC), BBSRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/04/2010		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes