# Influence of two local anaesthetic techniques on the survival of arteriovenous fistulas created for renal dialysis access

Submission date 27/11/2011	<b>Recruitment status</b> No longer recruiting	<ul> <li>Prospectively registered</li> <li>Protocol</li> </ul>
<b>Registration date</b> 18/01/2012	<b>Overall study status</b> Completed	<ul> <li>Statistical analysis plan</li> <li>Results</li> </ul>
Last Edited 27/03/2018	<b>Condition category</b> Nutritional, Metabolic, Endocrine	<ul> <li>Individual participant data</li> <li>Record updated in last year</li> </ul>

# Plain English summary of protocol

#### Background and study aims

Patients with terminal kidney disease (i.e. non-functioning kidneys) need to attend hospital for dialysis a few times a week, depending on how bad their kidney function is. The dialysis machine is attached to the patient by means of needles inserted in an arteriovenous fistula on the patient 's arm. An arteriovenous fistula provides easy access by means of a needle to a high flow of blood that is taken up the dialysis machine, processed, cleaned and returned back to the patient. These fistulas are created by sewing a vein onto an artery, usually in the forearm or elbow crease just below the skin. Once the fistula is ready it can be used time and time again for dialysis. The fistula operation can be done under general anaesthesia, regional anaesthesia (i.e. numbing the whole arm) or local anaesthesia (i.e. numbing the area where surgery takes place such as elbow crease or wrist). By far, most operations are done under local or regional anaesthesia with patients awake. The decision between local or regional anaesthesia is a matter of patient preference and local guidelines. Both techniques are widely accepted and used across the UK. Currently there is no evidence guiding best practice. Unfortunately for various reasons these fistulas tend to shut down over time. Studies have tried to address this but despite some advances, fistula failure still happens. At one year about 35% of fistulas fail, hence 3-4 out of 10 patients will require another fistula operation. Our study aims to compare the survival of fistulas created under regional anaesthesia versus local anaesthesia. There is recent evidence suggesting that creating these fistulas under regional anaesthesia helps to keep them open for longer. One of the main reasons this is thought to happen is because numbing the whole arm also causes the blood vessels to dilate (i.e. widen). This increases the flow of blood within those blood vessels and prevents clotting of the fistula. Wider blood vessels are also easier to see and handle and can make the operation easier and reduce blood vessel damage. Fistula operations are done with magnifying glasses so an increase in the size of blood vessels is always welcome.

Who can participate?

Patients aged 18 and over, who require the creation of a new arteriovenous fistula for dialysis.

### What does the study involve?

Participants are randomly allocated to one of two groups. Group A have their arteriovenous

fistula created by numbing the area where the surgery will take place (i.e. the elbow crease or the wrist). The numbing injection is given by the surgeon before the operation takes place. The numbness lasts 4-6 hours. Group B have their arteriovenous fistula created by numbing the whole arm. The numbing injection is given by an anaesthetist in the upper arm. They use an ultrasound machine to find the nerves of the arm and help them see exactly where to inject the local anaesthetic. The arm remains numb for up to 24 hours. Movement in that arm is also affected during that time. The fistula operation itself is the same in both groups. We are essentially comparing two anaesthetic approaches to fistula surgery. Both are safe, valid and used interchangeably at our hospital and throughout the country. We would like to know whether one is better.

What are the possible benefits and risks of participating?

By performing the fistula surgery after numbing the whole arm we hope to increase the survival of the fistula. Most fistula surgery in the UK is performed using local anaesthetic rather than general anaesthetic. There are very low risks of developing an allergy to the local anaesthetic. Regional anaesthesia of the arm is a very common and safe procedure, it is not experimental. It is a procedure performed on a daily basis at our hospital and throughout the country. The risks of having the fistula created by numbing the whole arm include injury to the nerves of the arm, which is a very rare complication that may happen to 1 patient in 10,000. The fact that an ultrasound machine is used to guide the injection reduces this risk. No such injuries have been recorded at our hospital despite daily practice. The arm will also be numb for up to 24 hours and movement will be affected. Some patients have reduced movement whilst others cannot use their am at all. The arm returns to normal after the local anaesthetic wears off; movement in the arm returns to normal earlier than sensation.

Where is the study run from? Sunderland Royal Hospital (UK)

When is the study starting and how long is it expected to run for? January 2012 to January 2014

Who is funding the study? Sunderland Royal Hospital (UK)

Who is the main contact? Dr Matei Dordea matei.dordea@nhs.net

# **Contact information**

**Type(s)** Scientific

**Contact name** Mr Andrew Brown

**Contact details** 

Sunderland Royal Hospital Department of Vascular Surgery Kayll Road Sunderland United Kingdom SR4 7TP +44 (0)191 565 6256 andrew.brown@chsft.nhs.uk

# Additional identifiers

EudraCT/CTIS number

**IRAS number** 

ClinicalTrials.gov number

Secondary identifying numbers N/A

# Study information

# Scientific Title

Influence of local and regional anaesthetic techniques on arteriovenous fistula access survival - a randomized controlled study

## Study objectives

There is an increasing amount of evidence that shows arteriovenous fistulas created under a regional block (i.e. axillary brachial plexus block hence anaesthetizing the whole arm) tend to remain patent for longer.

**Ethics approval required** Old ethics approval format

**Ethics approval(s)** NRES Committee North East Sunderland

**Study design** Randomised controlled trial

**Primary study design** Interventional

**Secondary study design** Randomised controlled trial

**Study setting(s)** Hospital

**Study type(s)** Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

## Health condition(s) or problem(s) studied

Surgical arteriovenous fistula for renal access dialysis

### Interventions

We are comparing two well established approaches to the creation of arteriovenous fistulas for renal access dialysis:

1. Arteriovenous fistula created under regional axillary block using levobupivacaine 2. Arteriovenous fistula created under local anaesthetic infiltration using levobupivacaine

Both groups will undergo Doppler assessment of their fistula at 8 weeks post-operatively to assess flow rates. This test is also undertaken by patients not taking part in the trial, hence not trial specific. Follow up for trial purposes is 8 weeks.

Please note this trial is not a Clinical Trial of Investigational Medicinal Products (CTIMP) (Medicines and Healthcare products Regulatory Agency [MHRA] guidance)

## Intervention Type

Drug

**Phase** Not Applicable

## Drug/device/biological/vaccine name(s)

Levobupivacaine

## Primary outcome measure

Fistula Doppler flow rate measurement at eight weeks post operatively. This measurement is already performed on all arteriovenous fistulas as part of routine arteriovenous fistula management.

## Secondary outcome measures

1. A patient satisfaction survey pertaining to patient experience of the local/regional anaesthetic techniques, the surgical procedure and their overall hospital experience will be undertaken

2. A standardized and validated questionnaire using a 5-point Likert scale will be provided to patients following the procedure and we will ask them to complete it before they leave hospital

# Overall study start date

01/01/2012

**Completion date** 01/01/2014

# Eligibility

Key inclusion criteria

 Patients referred for creation of a new radiocephalic or brachiocephalic fistula who consent to take part in the study and agree to have the procedure performed while conscious.
 Aged 18 - 90
 Male and female participants

### Participant type(s)

Patient

### Age group

Adult

## Lower age limit

18 Years

Sex

Both

# **Target number of participants** 220

## Key exclusion criteria

 Patients who require revision surgery.
 Having had previous arteriovenous fistula surgery is not a contraindication to recruitment provided the new procedure is not revision surgery
 Patients undergoing the procedure under General Anaesthetic

4. Known allergy to levobupivacaine (extremely rare)

## Date of first enrolment

01/01/2012

# Date of final enrolment

01/01/2014

# Locations

**Countries of recruitment** England

United Kingdom

**Study participating centre Sunderland Royal Hospital** Sunderland United Kingdom SR4 7TP

# Sponsor information

**Organisation** City Hospitals Sunderland NHS Foundation Trust (UK)

#### **Sponsor details**

c/o Ms Michelle Ferguson Directorate of General Surgery Department of Vascular Surgery Kayll Road Tyne & Wear Sunderland England United Kingdom SR4 7TP +44 (0)191 565 6256 michelle.ferguson@chsft.nhs.uk

**Sponsor type** Hospital/treatment centre

Website http://www.chsft.nhs.uk/

# Funder(s)

**Funder type** Hospital/treatment centre

**Funder Name** Sunderland Royal Hospital (UK)

# **Results and Publications**

**Publication and dissemination plan** Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

**IPD sharing plan summary** Not provided at time of registration