# Reducing Hyperpigmentation After Sclerotherapy

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>		
03/04/2020	No longer recruiting	☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
20/04/2020	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
11/12/2020	Skin and Connective Tissue Diseases			

#### Plain English summary of protocol

Background and study aims

Varicose veins are swollen blood vessels that are most commonly in the lower legs and feet. They may be dark blue or purple, and are often lumpy or twisted in appearance. People can experience swollen, itchy, aching and uncomfortable legs, feet and ankles. One treatment for varicose veins is called sclerotherapy, which involves an injection into the vein which causes the blood in the vein to clot and vein to collapse and shrink. One of the side effects of this treatment is that patients can be left with discolouration of the skin which can appear darker /hyperpigmented.

This study aims to find out if adding a drug called sulodexide to the standard sclerotherapy treatment for patients with varicose veins can reduce the presence of hyperpigmentation following the procedure without affecting the intended therapy of vein elimination or increasing the risk of major bleeding

#### Who can participate?

Patients who are aged 18 to 65 years with telangiectatic, reticular, or varicose veins in their lower limbs who are suitable candidates for sclerotherapy

#### What does the study involve?

Half of the participants will receive standard sclerotherapy treatment. The other half will receive sulodexide 7 days before the scheduled sclerotherapy. All participants will be advised to wear compression stockings for 7 days afterwards. At 1 and 3 months participants will return for follow up appointments to check for the success of the sclerotherapy and for hyperpigmentation.

What are the possible benefits and risks of participating?

Possible benefits of this treatment could be improved patient satisfaction, which is very important in a procedure mostly done for cosmetics reasons. Additionally, faster improvement in the emotional wellbeing and quality of life of the patient.

With the reduction of the incidence of post-sclerotherapy hyperpigmentation, the possibility to treat larger veins with sclerotherapy can be contemplated, and possibly avoid a more expensive

procedure like surgical stripping; this can be important in the socioeconomic population where sclerotherapy is the only possible treatment.

Where is the study run from? CLINEDEM, ISSSTECALI and Hospital General Issste 5 de Diciembre (Mexico)

When is the study starting and how long is it expected to run for? From January 2018 to September 2019

Who is funding the study? This study is investigator-initiated and funded

Who is the main contact?
Dr Alejandro Gonzalez Ochoa
2alex8as@gmail.com

## Contact information

#### Type(s)

Scientific

#### Contact name

Dr Alejandro Gonzalez Ochoa

#### **ORCID ID**

https://orcid.org/0000-0001-5068-623X

#### Contact details

Callejón 5 de Mayo and Calle 7 No. 791 Planta Alta San Luis Rio Colorado Mexico 83449 +526538497372 2alex8as@gmail.com

# Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

8936-042019

# Study information

Scientific Title

Reducing Hyperpigmentation After Sclerotherapy (RHyAS) study: a multicenter, randomized, clinical trial.

#### Acronym

RHyAS study

#### **Study objectives**

The use of sulodexide in patients with varicose vein treated with sclerotherapy can reduce the incidence of hyperpigmentation

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approved 21/11/2017, the Universidad Autónoma de Baja California (Coordination de Posgrado e Investigacion. av Alvaro Obregon y Julian Carrillo s/n Colonia Nueva cp 21100 Edificio de Rectoria, Mexicali Baja California, Mexico; +52 686 551 9497; anahernandez@uabcinvestigacion.net)

#### Study design

Prospective, multicentric, randomized controlled trial, using a parallel-group design

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Hyperpigmentation following sclerotherapy for varicose veins

#### **Interventions**

Participants who were included with telangiectasia, reticular or varicose veins who were candidates for sclerotherapy. They received Sclerotherapy using polidocanol 1% as sclerosant agent, 10ml maximum dose per session, total of 2 sessions 6 weeks apart. 20-30mmhg compression stockings were used in both groups for 7 days.

Participants were randomly assigned, in a 1:1 ratio, to either group A or group B.

Group A received an oral dose (250 LSU) of sulodexide bid 7 days prior to scheduled sclerotherapy that continued for 3 months.

Group B received the standard sclerotherapy protocol.

Photographic control was taken, and follow-up was done at 1 and 3 months. With the aid of computer software, the treated area was compared for the variables of incidence of pigmentation, the total area of pigmentation, skin-tone increase in pigmented area, vein disappearance, and incidence of major bleeding.

#### Intervention Type

Drug

#### Phase

Not Applicable

#### Drug/device/biological/vaccine name(s)

Suldexide 250 LSU bid oral dose for 3 months Polidocanol 1% 10ml maximum dose per session

#### Primary outcome(s)

- 1. Incidence of post sclerotherapy hyperpigmentation measured using computer software at baseline, 1 and 3 months
- 2. Total area of hyperpigmentation measured using computer software at baseline, 1 and 3 months

#### Key secondary outcome(s))

- 1. Presence of major bleeding measured using computer software at baseline, 1 and 3 months
- 2. Clinical response of vein disappearance measured using computer software at baseline, 1 and 3 months

#### Completion date

01/09/2019

# Eligibility

#### Key inclusion criteria

- 1. Aged 18 to 65 years
- 2. Telangiectatic, reticular, or varicose veins in lower limbs
- 3. Candidate for sclerotherapy
- 4. BMI 20 to 40 kg/m2
- 5. Fitzpatrick skin tone I V
- 6. Signed consent to participate

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Total final enrolment

720

#### Key exclusion criteria

- 1. Acute thrombosis
- 2. History of deep vein thrombosis
- 3. Deep vein reflux
- 4. Saphenous vein reflux
- 5. Pregnancy
- 6. Bed confinement
- 7. Long term use of steroidal or nonsteroidal anti-inflammatory drugs (such as corticoids, methotrexate, etc.)
- 8. Severe leg edema

#### Date of first enrolment

01/01/2018

#### Date of final enrolment

01/08/2019

## Locations

#### Countries of recruitment

Mexico

# Study participating centre

#### **CLINEDEM**

Clinic of Dental and Medical Specialties (CLINEDEM) Callejón 5 de Mayo and Calle 7 No. 791 Planta Alta San Luis Rio Colorado Mexico 83449

# Study participating centre ISSSTECALI

Av. Calafia No 1115 - 1G Centro Civico Mexicali Mexico 21000

#### Study participating centre Hospital General Issste 5 de Diciembre

Calzada independencia Centro Civico Mexicali Mexico 21000

# Sponsor information

#### Organisation

CLINEDEM

# Funder(s)

#### Funder type

Other

#### Funder Name

Investigator initiated and funded

# **Results and Publications**

#### Individual participant data (IPD) sharing plan

The datasets generated and/or analyzed during the current study during this study will be included in the subsequent results publication

#### IPD sharing plan summary

Data sharing statement to be made available at a later date

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2021	11/12/2020	Yes	No
Basic results			18/05/2020	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes