

# A pilot randomised controlled trial of antidepressant medication switching for treatment-refractory depression in primary care

<b>Submission date</b> 15/10/2004	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
<b>Registration date</b> 13/01/2005	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 28/10/2016	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr Paul Walters

### Contact details

MRC Fellow  
Section of Primary Care Mental Health  
Institute of Psychiatry  
Denmark Hill  
London  
United Kingdom  
SE5 8AF

-

p.walters@iop.kcl.ac.uk

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

## Secondary identifying numbers

PALBFWA

# Study information

## Scientific Title

A pilot randomised controlled trial of antidepressant medication switching for treatment-refractory depression in primary care

## Acronym

SWITCH

## Study objectives

The main hypothesis to be tested is:

In depressed primary care patients who have failed to recover after treatment with one antidepressant for at least four weeks and at a standard dose, switching their antidepressant medication to one from a different antidepressant class and continuing it for a further four weeks will be more effective than continuing the initial antidepressant for a further four weeks at the same dose.

Secondary hypotheses to be explored are:

1. That there is a positive association between social adversity, the number of life events and the response to antidepressant medications.
2. That non-compliance adversely affects the outcome independently of social adversity and life events
3. That both patient and doctors views of depression and antidepressant medication are associated with response to antidepressant medication

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Not specified

## Study type(s)

Treatment

## Participant information sheet

## **Health condition(s) or problem(s) studied**

Depression, Treatment-resistant

## **Interventions**

### **1. Antidepressant medication switched (intervention):**

Patients randomised to the intervention arm will have an appointment booked with their GP within 48 hours of the baseline assessment. At this appointment the GP will follow pre-determined guidelines for switching the Selective Serotonin Reuptake Inhibitor (SSRI) to an alternative antidepressant from a different antidepressant class. The Maudsley Prescribing Guidelines (6th Edition) will be used to guide General Practitioners (GPs) switching antidepressants (incorporating the recommended wash-out periods between antidepressant medications). Patients will continue on the second antidepressant for four weeks at a standard dose.

For patients unable to tolerate the second antidepressant due to side effects, GPs will follow guidelines to prescribe an alternative antidepressant so that each patient will have a full trial of four weeks on a second antidepressant from a different antidepressant class. The GP will be asked to assess the patient at two and four weeks (following best-practice guidelines) but can see the patient more often as deemed necessary and refer for counselling as necessary.

### **2. Same antidepressant medication continued (control):**

Patients randomised to the control arm will have an appointment booked with their GP within 48 hours of the baseline assessment. Patients randomised to the control group will be asked by their GP to continue on the initial SSRI for a further four weeks. The GP will be asked to assess the patient at two and four weeks (following best-practice guidelines) but can see the patient more often as deemed necessary and refer for counselling as necessary.

## **Intervention Type**

Other

## **Phase**

Not Specified

## **Primary outcome measure**

To determine the feasibility of conducting a larger randomised controlled trial to examine clinical effectiveness of switching an antidepressant medication to another from a different antidepressant class in primary care patients with treatment-refractory depression.

## **Secondary outcome measures**

1. Whether demographic variables, social adversity, life events, physical health and disability, are associated with outcome
2. Whether patient and doctor beliefs about depression and antidepressant medication are associated with outcome

## **Overall study start date**

01/02/2005

## **Completion date**

30/09/2006

## **Eligibility**

**Key inclusion criteria**

Consenting patients aged 16 years or over with a score of 13 or more on the Hamilton Rating Scale for Depression (HAM-D), 9 or more on the Beck Depression Inventory and who have been compliant with a specific serotonin reuptake inhibitor at a standard dose for at least four weeks (as ascertained by self report questionnaire) will be invited to participate in the study.

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

100

**Key exclusion criteria**

1. Temporary registrations
2. Under 16 years
3. Currently receiving psychological therapies (cognitive behavioural therapy, problem solving therapy, interpersonal therapy)
4. Other psychotropic medications apart from benzodiazepines
5. Case note diagnosis of schizophrenia or bipolar affective disorder
6. Currently being treated for drug or alcohol dependence
7. Suicidal patients
8. Psychotic depression as evidenced by delusions and/or hallucinations
9. Patient treated by secondary psychiatric services
10. Non-English speaking

**Date of first enrolment**

01/02/2005

**Date of final enrolment**

30/09/2006

**Locations****Countries of recruitment**

England

United Kingdom

**Study participating centre**

**MRC Fellow**  
London  
United Kingdom  
SE5 8AF

## Sponsor information

**Organisation**  
King's College London (UK)

**Sponsor details**  
Institute of Psychiatry  
De Crespigny Park  
London  
England  
United Kingdom  
SE5 8AF  
-  
G.Dale@iop.kcl.ac.uk

**Sponsor type**  
University/education

**ROR**  
<https://ror.org/0220mzb33>

## Funder(s)

**Funder type**  
Research council

**Funder Name**  
MRC Clinical research training fellowship

## Results and Publications

**Publication and dissemination plan**  
Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**  
Not provided at time of registration