Comparison of efficacy and safety of two forms of cyclodestructive therapy. Transscleral cyclophotocoagulation and endoscopic cyclophotocoagulation in the treatment of patients with refractory glaucoma.

| Submission date | Recruitment status No longer recruiting | Prospectively registered | | |
|-------------------------------|---|--|--|--|
| 12/09/2003 | | ☐ Protocol | | |
| Registration date | Overall study status | Statistical analysis plan | | |
| 12/09/2003 | Completed | [X] Results | | |
| Last Edited 06/01/2009 | Condition category Eye Diseases | [] Individual participant data | | |

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Mr Philip Bloom

Contact details

Ophthamology Western Eye Hospital Marylebone Road London United Kingdom NW1 5YE +44 (0)20 7886 6666

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N0241032308

Study information

Scientific Title

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Not Specified

Participant information sheet

Not provided at time of registration

Health condition(s) or problem(s) studied

Eye Diseases: Glaucoma

Interventions

Transscleral cyclophotocoagulation vs endoscopic cyclophotocoagulation

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Not provided at time of registration

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/11/1998

Completion date

01/07/2005

Eligibility

Key inclusion criteria

100 Patients will be recruited, randomised into two groups.

Participant type(s)

Patient

Age group

Not Specified

Sex

Not Specified

Target number of participants

100 patients will be recruited, randomised into two groups.

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/11/1998

Date of final enrolment

01/07/2005

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

Ophthamology

London United Kingdom NW1 5YE

Sponsor information

Organisation

Department of Health (UK)

Sponsor details

Richmond House 79 Whitehall London United Kingdom SW1A 2NL

Sponsor type

Government

Website

http://www.doh.gov.uk

Funder(s)

Funder type

Government

Funder Name

St Mary's NHS Trust (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-----------------|---------|--------------|------------|----------------|-----------------|
| Results article | results | 01/06/2006 | | Yes | No |