

The effect of timing of the application of topical anaesthetic prior to rigid nasendoscopy

Submission date 29/09/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 29/09/2006	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 08/12/2010	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr David Pothier

Contact details
ENT
Royal United Hospital
B&NES
Bath
United Kingdom
BA1 3NG
+44 (0)1225 428331
David.Pothier@ruh-bath.swest.nhs.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0212177261

Study information

Scientific Title

Study objectives

Is it best to wait one or ten minutes before performing nasendoscopy after the application of cophenylcaine?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Not Specified

Participant information sheet

Health condition(s) or problem(s) studied

Nasendoscopy

Interventions

Quantitative, randomised controlled trial

Treatment 1: wait 1 minute

Treatment 2: wait 10 minutes

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome measure

Visual analogue scales of pain and discomfort for patients.

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/03/2006

Completion date

01/09/2006

Eligibility

Key inclusion criteria

Patients who, by the decision of the examining clinician, need to undergo examination of their nasal cavity using rigid nasendoscopy.

Participant type(s)

Patient

Age group

Not Specified

Sex

Not Specified

Target number of participants

50

Key exclusion criteria

1. Unable to provide informed consent
2. Patients in whom lignocaine or phenylphrine is contra indicated for medical reasons
3. Patients in whom only one nasal cavity or part of one cavity is to be examined

Date of first enrolment

01/03/2006

Date of final enrolment

01/09/2006

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

ENT
Bath
United Kingdom
BA1 3NG

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2006 Update - Department of Health

Sponsor details

The Department of Health, Richmond House, 79 Whitehall
London
United Kingdom
SW1A 2NL
+44 (0)20 7307 2622
dhmail@doh.gsi.org.uk

Sponsor type

Government

Website

<http://www.dh.gov.uk/Home/fs/en>

Funder(s)

Funder type

Government

Funder Name

Royal United Hospital Bath NHS Trust

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2007		Yes	No