

A randomised controlled trial of cognitive therapy for command hallucinations

Submission date

23/01/2004

Recruitment status

No longer recruiting

☐ Prospectively registered

☐ Protocol

Registration date

23/01/2004

Overall study status

Completed

☐ Statistical analysis plan

☒ Results

Last Edited

29/01/2010

Condition category

Mental and Behavioural Disorders

☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Peter Trower

Contact details

University of Birmingham / South Bham Mental Health Trust
Queen Elizabeth Psychiatric Hospital
Mindelsohn Way
Birmingham
United Kingdom
B15 2TT

Additional identifiers

Protocol serial number

RGC00391

Study information

Scientific Title**Study objectives**

The objective is to evaluate whether a form of cognitive therapy (CT) developed by the applicants is an effective form of treatment for command hallucinations. The researcher's cognitive model asserts that high risk behaviours of command hallucinators are a consequence of core beliefs, namely a conviction in the power of their voices and a belief that they are compelled to comply with and appease these voices.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Schizophrenia and other psychoses

Interventions

1. Cognitive therapy
2. Control

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. Voices:
 - 1.1. Content - Cognitive Assessment Schedule (Chadwick & Birchwood, 1995)
 - 1.2. Topograph - Frequency, volume, audibility, affect (Hustig & Hafner, 1992)
 - 1.3. Beliefs and evidence
 - 1.3.1. Beliefs About Voices Questionnaire (Chadwick & Birchwood, 1995)
 - 1.3.2. Maudsley Assessment of Delusions Schedule (Buchanan et al 1993)
 - 1.4. Compliance (Beck-Sander et al 1997)
2. Affect:
 - 2.1. Depression, hopelessness (Calgary Depression Scale for Schizophrenia: Addington & Addington, 1993)
3. Mental State:
 - 3.1. Positive and Negative Syndrome Scale (PANSS) (Kay, 1997)
 - 3.2. Previous episodes of 'risk' behaviours

Key secondary outcome(s)

Not provided at time of registration

Completion date

01/10/2001

Eligibility

Key inclusion criteria

The sample will be drawn from the case registers of patients receiving treatment from South and North Birmingham Mental Health Trusts. From this sample base the researchers will draw a cohort of patients conforming to International Statistical Classification of Diseases and Related Health Problems, Tenth revision (ICD-10) schizophrenia who have experienced auditory command hallucinations for a minimum of six months. There will be 26 patients in each of the groups.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Current drug and alcohol misuses
2. Primary organic disorder

This is to ensure that the auditory hallucinations are a symptom of schizophrenia rather than induced by drugs or the result of organic impairment.

Date of first enrolment

01/10/1999

Date of final enrolment

01/10/2001

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

University of Birmingham / South Bham Mental Health Trust
Birmingham
United Kingdom
B15 2TT

Sponsor information

Organisation

NHS R&D Regional Programme Register - Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

NHS Executive West Midlands (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/04/2004		Yes	No