

# Effect of high volume saline enemas during labour

<b>Submission date</b> 20/03/2002	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 20/03/2002	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 04/10/2017	<b>Condition category</b> Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Luis Gabriel Cuervo

**Contact details**  
525 23rd St, NW  
Washington  
United States of America  
DC 20037-2895  
+ 1 202 974 3135  
lgcuervo@gmail.com

## Additional identifiers

**Protocol serial number**  
N/A

## Study information

**Scientific Title**  
Effect of high volume saline enemas during labour: a randomised controlled trial

**Study objectives**

The main objective of this trial was to determine if the use of high volume enemas during the first stage of labour modified neonatal and/or puerperal infectious rates. The null hypothesis stated that infectious morbidity was similar for intervention and control groups. A secondary objective was to establish if there is any effect on specific neonatal or puerperal infectious rates and other clinically relevant outcomes.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Randomised controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Puerperal and neonatal infections

**Interventions**

Enema versus no enema. Block randomisation allocation with sealed envelopes.

**Intervention Type**

Other

**Phase**

Not Specified

**Primary outcome(s)**

The primary outcomes were infections in newborns and /or puerperal women. During the analysis, neonatal and puerperal infections were described individually and then amalgamated into a new outcome. This outcome was the 'combined maternal-neonatal outcome'.

A neonatal outcome was positive when during follow-up, any of the following clinical conditions were diagnosed : ocular infection (defined as purulent drainage in the eye after the sixth day of delivery); umbilical infection (foul smell with periumbilical erythema); skin infection (cellulitis or impetigo); respiratory tract infection (clinical diagnosis of lower or upper respiratory tract infection); intestinal infection; meningitis, sepsis or if the child had been prescribed systemic antibiotics during the first month of life.

Positive puerperal outcomes were registered when the mother had any of the following conditions diagnosed by a health care provider: any suture dehiscence, purulent effusion from the episiorraphy, urinary tract infection, pelvic inflammatory disease or vulvovaginitis. While hospitalised, participating women and newborns were visited on a daily basis. Throughout the process data were gathered by trained research assistants using standardised questionnaires.

Formats were also used to register data retrieved from telephone calls, during hospitalisation, at follow-up visits and when any communication was established with participants, their families or health care providers.

**Key secondary outcome(s)**

Not provided at time of registration

**Completion date**

28/02/1998

## **Eligibility**

**Key inclusion criteria**

1. Women attending labour ward at low risk for delivery
2. Greater than 36 week gestation
3. In labour
4. Cervix dilatation <8 cm

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

Female

**Key exclusion criteria**

Clinical diagnosis of any systemic or gynaecological bacterial infection, use of systemic antibiotics during the week prior to admission, rupture of amniotic membranes or doubt of their integrity, and a cervical dilatation greater than or equal to 7 cm.

**Date of first enrolment**

01/02/1997

**Date of final enrolment**

28/02/1998

## **Locations**

**Countries of recruitment**

Colombia

United States of America

**Study participating centre**  
525 23rd St, NW  
Washington  
United States of America  
DC 20037-2895

## Sponsor information

**Organisation**  
INCLEN Trust (USA)

## Funder(s)

**Funder type**  
Research organisation

**Funder Name**  
International Clinical Epidemiology Network seed grant

**Funder Name**  
Enemas donated by Baxter (without detailed knowledge of research protocol)

**Funder Name**  
Own funds provided by Luis Gabriel Cuervo

**Funder Name**  
Logistics by the School of Medicine at the Universidad Javeriana in Bogota

## Results and Publications

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**  
Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	Results	19/03/2006		Yes	No