

# A comparison of intermittent aortic cross clamping and intermittent aortic warm blood cardioplegia for myocardial protection during coronary artery bypass surgery

<b>Submission date</b>	<b>Recruitment status</b>	<input type="checkbox"/> Prospectively registered
30/09/2005	No longer recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
30/09/2005	Completed	<input type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
13/03/2014	Surgery	<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Mr JAJ Hyde

### Contact details

Cardiac Services

Queen Elizabeth Hospital

Birmingham

United Kingdom

B15 2TH

## Additional identifiers

### Protocol serial number

N0265006856

## Study information

### Scientific Title

## **Study objectives**

Is the newer technique of intra-aortic warm blood cardioplegia superior to the established technique of intermittent aortic cross-clamping for myocardial protection during coronary surgery?

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Not provided at time of registration

## **Study design**

Randomised controlled trial

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Surgery: Coronary artery bypass grafting (CABG)

## **Interventions**

40 patients undergoing elective CABG will randomly be allocated into two groups of 20 patients each, to receive either intermittent cross clamp with fibrillation (XCF) or intermittent warm blood cardioplegia (WBC). Blood will be taken from in-situ intra-arterial or intravenous cannulae, at the following 12 time points: Following induction of anaesthesia; Following institution of cardiopulmonary bypass, but prior to the application of an aortic cross clamp with resultant myocardial ischaemia; 30,60,90,120 minutes and 4,6,8,12,24,28 hours following final release of the aortic cross clamp. The blood will be centrifuged for serum, which will be stored frozen at -70 C until batch analysis. The analysis will be performed blind.

## **Intervention Type**

Procedure/Surgery

## **Phase**

Not Specified

## **Primary outcome(s)**

Not provided at time of registration

## **Key secondary outcome(s)**

Not provided at time of registration

## **Completion date**

01/01/2007

## **Eligibility**

**Key inclusion criteria**

Patients undergoing elective CABG

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Not Specified

**Sex**

Not Specified

**Key exclusion criteria**

Not provided at time of registration

**Date of first enrolment**

01/01/2004

**Date of final enrolment**

01/01/2007

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre**

Cardiac Services

Birmingham

United Kingdom

B15 2TH

## Sponsor information

**Organisation**

Department of Health

## Funder(s)

**Funder type**

Government

**Funder Name**

University Hospital Birmingham NHS Trust (UK)

**Funder Name**

Research Funds

## Results and Publications

### Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not provided at time of registration