# Treatment with medications that prevent the formation of blood clots, in the primary prevention of heart disease in patients with a stable lung condition called Chronic Obstructive Pulmonary Disease (COPD)

Submission date	Recruitment status No longer recruiting	<ul><li>Prospectively registered</li></ul>		
21/10/2015		[X] Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
21/10/2015		[X] Results		
<b>Last Edited</b> 08/11/2023	Condition category Respiratory	[] Individual participant data		

### Plain English summary of protocol

Background and study aims

Chronic obstructive pulmonary disease (COPD) is the collective name for a group of diseases which affect the lungs. People who suffer from COPD have difficulty breathing, which gets worse over time. This is because the airways become narrowed or blocked, and the air sacs (alveoli) in the lungs are destroyed or lose their ability to stretch. The main cause of COPD is smoking, which over time permanently damages the lining of the lungs. Many studies have shown that people with COPD have a greater risk of developing heart and blood vessel disease (cardiovascular disease) which can lead to a heart attack. When a person is suffering from cardiovascular disease (CVD), fatty substances (plaque) build up inside blood vessels causing them to become narrowed. Platelets (a component of blood) "stick" to the walls of these narrowed blood vessels, causing blockages (occlusion) that can lead to a heart attack. Ticagrelor is a medication which works by slowing and stopping platelets from sticking to the blood vessel walls. It is usually taken with aspirin, which makes this more effective by "thinning" the blood. These medications are often used in CVD patients and have been shown to be very effective. Currently there is very little information about the best way to treat COPD patients who are at risk of heart disease. The aim of this study is to find out whether treatment with ticagrelor and /or aspirin can help to lower the risk of heart disease in COPD patients.

### Who can participate?

Adults who have smoked for at least 10 years with signs of COPD.

### What does the study involve?

Participants are randomly allocated to one of three treatment groups or a control group. Those in the treatment groups are given aspirin and/or ticagrelor to take, and those in the control group are given a placebo (dummy pill). At the initial visit, participants will undergo a blood test, a lung function test, an ultrasound scan, they will have their stiffness of the blood vessels

measured and they will be asked to complete questionnaires. Participants who are receiving the treatment attend follow up visits at 1, 3 and 6 months to see whether the medication is having any effect. The blood tests and questionnaires are repeated at a 6 month follow up visit for both groups.

What are the possible benefits and risks of participating?

There are no direct benefits to the patient for participating in this study. However, their participation in this study may add to the medical knowledge about how to treat COPD patients who are at risk of future heart problems. There are minor risks related to taking the study medications (asprin and ticagrelor), however the benefits have been judged to outweigh the risks.

Where is the study run from? Newcastle upon Tyne Hospitals NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for? September 2015 to November 2017

Who is funding the study? AstraZeneca UK Limited (UK)

Who is the main contact? Catherine Brennand cath.brennand@ncl.ac.uk

## Contact information

### Type(s)

Public

### Contact name

Ms Catherine Brennand

### Contact details

Newcastle Clinical Trials Unit Faculty of Medical Sciences 1-4 Claremont Terrace Newcastle University Newcastle Upon Tyne United Kingdom NE2 4AE +44 (0)191 208 7258 cath.brennand@ncl.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS) 2014-005475-86

ClinicalTrials.gov (NCT)

### Protocol serial number

**CPMS 19244** 

# Study information

### Scientific Title

Anti-platelet therapy in the primary prevention of cardiovascular disease in patients with chronic obstructive pulmonary disease

### Acronym

APPLE-COPD:ICON 2

### Study objectives

The aim of this study is to find out whether ticagrelor and/or aspirin are effective treatments in the primary prevention of cardiovascular disease in patients with chronic obstructive pulmonary disease.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

First Medical Research Ethics Commitee, 26/05/2015, ref: 15/NE/0155

### Study design

Randomized; Interventional; Design type: Treatment

### Primary study design

Interventional

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Topic: Cardiovascular disease, Respiratory disorders; Subtopic: Cardiovascular (all Subtopics), Respiratory (all Subtopics); Disease: Cardiovascular, Respiratory

### Interventions

Patients who meet the inclusion criteria with a QRISK2 score >20% will be randomised to the treatment arm. A stratified blocked treatment allocation system is used to randomise patients into four groups who each receive a different treatment for six months:

- 1. Aspirin (75 mg once daily dose) and Ticagrelor (90 mg twice daily dose)
- 2. Aspirin (75 mg once daily dose) and placebo
- 3. Ticagrelor (90 mg twice daily dose) and placebo
- 4. Placebo alone

Block size will not be disclosed to the investigators. Randomisation will be administered centrally via the Newcastle Clinical Trials Unit using a secure web-based system.

Patients will attend follow-up visits at the Clinical Research Facility, Royal Victoria Infirmary at 1-month, 3-months and 6-months and then followed up clinically for up to 1 year.

Patients with a QRISK2 score <20% will be allocated to the observational cohort of the study, they will undergo the baseline procedures, including blood tests and will be followed up again at 1 year. Patients in the observational cohort will not undergo follow-up visits at 1 month, 3 months and 6 months.

### Intervention Type

Drug

### Phase

Phase II

### Drug/device/biological/vaccine name(s)

Aspirin, ticagrelor

### Primary outcome(s)

Inhibition of adenosine disphosphate-induced platelet aggregation measured using the multiplate test at baseline and 6 months.

### Key secondary outcome(s))

- 1. Changes in inflammatory markers including fibrinogen, hsCRP, TNF alpha, IL-6, MPO is measured from blood samples taken at baseline and 6 months
- 2. Changes in carotid intima media thickness measured using an ultrasound probe and vascular stiffness measured using a specialised pressure cuff similar to a blood pressure cuff at baseline and 6 months
- 3. Quality of life measured using questionnaires (EQ5D-5L, St. George's COPD questionnaire) at baseline and 6 months

### Completion date

10/11/2017

# Eligibility

### Key inclusion criteria

- 1. Aged 18 years or over
- 2. Abnormal spirometry with FEV1<80% and FEV1/FVC ratio <70% of predicted
- 3. Smoking history that is 10 years or greater (current or ex smokers can be included)
- 4. Patient has the capacity to consent

### Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Adult

### Lower age limit

18 years

### Sex

All

### Total final enrolment

120

### Key exclusion criteria

- 1. Any condition that is being concurrently treated through anticoagulation or antiplatelet therapy (atrial fibrillation, deep vein thrombosis, valve prosthesis, recent myocardial infarction, use of drug eluting stents)
- 2. Other specific contraindications to management with antiplatelet medication (bleeding risks, allergies)
- 3. Any contraindication for Aspirin and Ticagrelor use
- 4. Other concurrent terminal illnesses with life expectancy less than 1 year (congestive cardiac failure, carcinoma etc.)
- 5. Current involvement in another clinical trial or exposure to another IMP within the previous 30 days
- 6. COPD with an atypical cause (e.g. A1antitrypsin deficiency)
- 7. Planned/ Expected major surgery where anti-platelet therapy would be ceased
- 8. Pregnancy, planned pregnancy or current breastfeeding
- 9. Patient is unable to provide informed consent
- 10. Younger than 18 years

### Date of first enrolment

04/09/2015

### Date of final enrolment

31/05/2017

### Locations

### Countries of recruitment

United Kingdom

England

# Study participating centre Royal Victoria Infirmary

Newcastle upon Tyne Hospitals NHS Foundation Trust Queen Victoria Road Newcastle Upon Tyne United Kingdom NE1 4LP

### Study participating centre

### Freeman Hospital

Newcastle upon Tyne Hospitals NHS Foundation Trust Freeman Road Newcastle Upon Tyne United Kingdom NE7 7DN

# Sponsor information

### Organisation

Newcastle upon Tyne Hospitals NHS Foundation Trust

### ROR

https://ror.org/05p40t847

# Funder(s)

### Funder type

Industry

### **Funder Name**

AstraZeneca

### Alternative Name(s)

AstraZeneca PLC, Pearl Therapeutics, AZ

### **Funding Body Type**

Government organisation

### **Funding Body Subtype**

For-profit companies (industry)

### Location

United Kingdom

### **Results and Publications**

### Individual participant data (IPD) sharing plan

There currently are no provisions or a plan in place for data sharing for APPLE-COPD. The data are held by NCTU on behalf of the Sponsor. NCTU are currently developing a data sharing policy and it may be possible to make data available in the future.

**IPD sharing plan summary**Not expected to be made available

# Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		05/08/2019	08/11/2023	Yes	No
Protocol article	protocol	26/05/2018	17/05/2019	Yes	No
Basic results			17/06/2020	No	No
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes