

TyVAC Nepal: typhoid vaccine study

| | | |
|----------------------------------------|----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|
| Submission date 20/06/2017 | Recruitment status No longer recruiting | <input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol |
| Registration date 26/09/2017 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results |
| Last Edited 17/10/2025 | Condition category Infections and Infestations | <input type="checkbox"/> Individual participant data |

Plain English summary of protocol

Current plain English summary as of 13/08/2018:

Background and study aims

Typhoid fever is a bacterial infection that can cause severe disease and even death. Typhoid is spread through contaminated food and water, due to poor hygiene and sanitation conditions. In Nepal, typhoid fever causes a lot of disease, mostly in children. Previous vaccines have not provided long-term protection for children. A new typhoid vaccine, Vi-TCV, has been developed and is licensed for use in Nepal. This vaccine is safe for children and may provide long-term protection. However, this vaccine is not yet available through the routine childhood immunisation programme. Before the government will introduce the vaccine into the routine programme, more information is needed on the level of protection that the vaccine provides. The aim of this study is to find out whether the typhoid vaccine Vi-TCV reduces the incidence of typhoid fever in children.

Who can participate?

Children under the age of 16

What does the study involve?

Participants are randomly allocated to either receive the typhoid vaccine (Vi-TCV) or a vaccine which protects against meningitis. Participants are followed-up over a two-year period. When participants develop a fever during that time, they are asked to come to the local hospital or the study clinics in the community. If they have a fever over 38 degrees C or if they have had a fever for 2 or more days they receive treatment and a blood test to check if they have typhoid fever. The cost of treatment for suspected or confirmed typhoid is covered by the study. All participants are also monitored for side effects associated with the vaccine. At the end of the study, all of the children who first received the meningitis vaccine are offered the typhoid vaccine too.

What are the possible benefits and risks of participating?

Participants benefit from the study by receiving the typhoid vaccine for free, which is not currently available through the routine immunisation system in Nepal. Participants also have free treatment for any suspected or confirmed typhoid infections. The vaccine has been found to be safe in all previous studies, but since this is the largest study of this vaccine to date, there is a risk for participants that a rare side effect, not yet identified, may become apparent in this study.

Where is the study run from?
Patan Hospital (Nepal)

When is the study starting and how long is it expected to run for?
November 2017 to August 2020

Who is funding the study?
Bill and Melinda Gates Foundation (USA), grant number OPP1151153

Who is the main contact?
Prof. Andrew Pollard

Previous plain English summary:

Background and study aims

Typhoid fever is a bacterial infection that can cause severe disease and even death. Typhoid is spread through contaminated food and water, due to poor hygiene and sanitation conditions. In Nepal, typhoid fever causes a lot of disease, mostly in children. Previous vaccines have not provided long-term protection for children. A new typhoid vaccine, Vi-TCV, has been developed and is licensed for use in Nepal. This vaccine is safe for children and may provide long-term protection. However, this vaccine is not yet available through the routine childhood immunisation programme. Before the government will introduce the vaccine into the routine programme, more information is needed on the level of protection that the vaccine provides. The aim of this study is to find out whether the typhoid vaccine Vi-TCV reduces the incidence of typhoid fever in children.

Who can participate?
Children under the age of 16

What does the study involve?

Participants are randomly allocated to either receive the typhoid vaccine (Vi-TCV) or a vaccine which protects against meningitis. Participants are followed-up over a two-year period. When participants develop a fever during that time, they are asked to come to the local hospital or the study clinics in the community. If they have a fever over 38 degrees C or if they have had a fever for 2 or more days they receive treatment and a blood test to check if they have typhoid fever. The cost of treatment for suspected or confirmed typhoid is covered by the study. All participants are also monitored for side effects associated with the vaccine. At the end of the study, all of the children who first received the meningitis vaccine are offered the typhoid vaccine too.

What are the possible benefits and risks of participating?

Participants benefit from the study by receiving the typhoid vaccine for free, which is not currently available through the routine immunisation system in Nepal. Participants also have free treatment for any suspected or confirmed typhoid infections. The vaccine has been found to be safe in all previous studies, but since this is the largest study of this vaccine to date, there is a risk for participants that a rare side effect, not yet identified, may become apparent in this study.

Where is the study run from?
Patan Hospital (Nepal)

When is the study starting and how long is it expected to run for?
November 2017 to August 2020

Who is funding the study?
Bill and Melinda Gates Foundation (USA)

Who is the main contact?
Prof. Andrew Pollard

Contact information

Type(s)
Scientific

Contact name
Prof Andrew Pollard

ORCID ID
<https://orcid.org/0000-0001-7361-719X>

Contact details
Oxford Vaccine Group
Centre for Clinical Vaccinology and Tropical Medicine (CCVTM)
Churchill Hospital Old Road
Oxford
United Kingdom
OX3 7LE

Additional identifiers

Protocol serial number
OVG2017/05

Study information

Scientific Title
Assessing the impact of a Vi-Polysaccharide Conjugate Vaccine in preventing typhoid infection among Nepalese children – a Phase III trial

Study objectives
The typhoid vaccine, Vi-TCV, will reduce the incidence of laboratory confirmed typhoid fever in children receiving the vaccine, compared to those receiving a control vaccine.

Ethics approval required
Old ethics approval format

Ethics approval(s)
1. Nepal Health Research Council (NHRC), 16/08/2017, reference number: 386, registration number: 170/2017
2. Oxford Tropical Research Ethics Council (OxTREC), 10/08/2017, ref: 15-17

Study design

Participant- and observer-blind randomised-controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Blood culture confirmed typhoid fever

Interventions

Current interventions as of 13/08/2018:

Participants will be randomised (block randomisation) 1:1 to receive a single dose of either Vi-TCV or MenA vaccine:

Intervention: Vi Typhoid conjugate vaccine (Vi-TCV), trade name: TyBar

Control: Meningococcal group A vaccine (MenA), trade name: MenAfriVac

Neither the participants or the researchers will know which vaccine any child has received until the end of trial. At the end of the trial, once it is unblinded, and everyone knows which vaccine they received, all of the children who first received the control vaccine will be offered the typhoid vaccine too, and all children who received the intervention vaccine will also be offered the control meningococcal group A vaccine.

Participants will be followed-up over a two year period. When participants develop a fever during that time, they will be asked to come to the local hospital or the trial clinics in the community. If they have a fever over 38 degrees C, or if they have had a fever for 2 or more days, they will get treatment and a blood test to check if they have typhoid fever. The cost of treatment for suspected or confirmed typhoid will be covered by the trial. All participants will also be monitored for safety outcomes associated with the vaccine.

Previous interventions:

Participants will be randomised (block randomisation) 1:1 to receive a single dose of either Vi-TCV or MenA vaccine:

Intervention: Vi Typhoid conjugate vaccine (Vi-TCV), trade name: TyBar

Control: Meningococcal group A vaccine (MenA), trade name: MenAfriVac

Neither the participants or the researchers will know which vaccine any child has received until the end of trial. At the end of the trial, once it is unblinded, and everyone knows which vaccine they received, all of the children who first received the control vaccine will be offered the typhoid vaccine too.

Participants will be followed-up over a two year period. When participants develop a fever during that time, they will be asked to come to the local hospital or the trial clinics in the community. If they have a fever over 38 degrees C, or if they have had a fever for 2 or more days, they will get treatment and a blood test to check if they have typhoid fever. The cost of treatment for suspected or confirmed typhoid will be covered by the trial. All participants will also be monitored for safety outcomes associated with the vaccine.

Intervention Type

Biological/Vaccine

Phase

Phase III

Drug/device/biological/vaccine name(s)

TyBar (Vi-TCV), MenAfriVac

Primary outcome(s)

The efficacy and rate reduction of the Vi-TCV in preventing blood culture-confirmed symptomatic infection caused by *S. Typhi*, measured through the incidence of blood culture confirmed typhoid fever in participants presenting to hospital or clinic for the duration of the trial, in each vaccination arm

Key secondary outcome(s)

1. Vi-TCV vaccine safety, measured as the proportion of participants developing all adverse events within the first 7 days post-vaccination, and serious adverse events within 6 months of vaccination, as determined through self-reporting at three-monthly follow-up contact
2. The impact of vaccination with Vi-TCV on the incidence of inpatient/outpatient admission rates for fever, measured by the rates of participants with at least ≥ 2 days of subjective persistent fever, or a temperature of at least 38 degrees C, at presentation at Patan Hospital or trial clinics in each vaccination arm, stratified by duration and severity of fever
3. The difference in rates of hospital and clinic presentation for febrile illness in each vaccination arm, measured as the rates of hospital or clinic presentation with febrile illness of any duration in each vaccination arm, measured by hospital presentation logs, hospital records, trial clinic records and self-reporting during three monthly follow-up
4. Days spent in hospital from febrile illness, measured as the length of stay in hospital, collected from Patan hospital patient records, and parent/self-reported, in each vaccination arm
5. Incidence of clinically-suspected enteric fever, measured as the number of clinical diagnoses of typhoid fever, as determined by trial staff in Patan hospital outpatient clinics and trial clinic, in each vaccination arm throughout the two year follow-up period
6. Paratyphoid infection rates in each vaccination arm, measured as the rates of blood culture confirmed Paratyphoid cases in participants presenting to hospital or clinic for the duration of the trial, in each vaccination arm

Added 20/06/2019:

7. The efficacy and rate reduction of the Vi-TCV in preventing blood culture-confirmed symptomatic infection caused by *S. Typhi*, measured through the incidence of blood culture confirmed typhoid fever in participants presenting to hospital or clinic with at least 3 days of fever, for the duration of the trial in each vaccination arm

Completion date

01/08/2020

Eligibility

Key inclusion criteria

1. Parent/legal guardian is willing and competent to provide informed consent. If the participant is 12 years of age or older, informed assent will also be sought
2. Aged between 9 months (or eligible for measles vaccination according to local protocol) and <16 years (i.e. up to 15 years 364 days) at time of vaccination
3. In good health on the day of vaccination

4. Parent/legal guardian confirms that their child will be willing and be able to comply with study requirements including follow-up contact, according to the trial schedule
5. Live within the study catchment area at the time of vaccination

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Child

Lower age limit

9 months

Upper age limit

16 years

Sex

All

Total final enrolment

20019

Key exclusion criteria

The participant will not be enrolled if any of the following criteria apply:

1. They have knowingly received a typhoid vaccine in the last three years
2. They have a known allergy to any of the vaccine components
3. Any medical or social reasons that will prevent the participant from conforming to the study requirements as judged by a medical professional
4. They are planning to move away from the catchment area within the next 6 months

Participants will be temporarily excluded from being vaccinated if, at point of vaccination, any of the following apply:

1. Reported fever within 24 hours prior to vaccination
2. Use of anti-pyretics within 4 hours prior to vaccination

If these apply, the participant will be temporarily excluded for vaccination until 48 hours has passed. A re-assessment will be conducted to ensure these temporary exclusion criteria no longer exist

Date of first enrolment

01/11/2017

Date of final enrolment

09/04/2018

Locations**Countries of recruitment**

Nepal

Study participating centre

Patan Hospital

Kathmandu

Nepal

-

Sponsor information

Organisation

University of Oxford

ROR

<https://ror.org/052gg0110>

Funder(s)

Funder type

Charity

Funder Name

Bill and Melinda Gates Foundation

Alternative Name(s)

Bill & Melinda Gates Foundation, Gates Foundation, Gates Learning Foundation, William H. Gates Foundation, BMGF, B&MGF, GF

Funding Body Type

Government organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United States of America

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Prof. Andrew Pollard.

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-----------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|--------------|------------|----------------|-----------------|
| Results article | results | 05/12/2019 | 06/12/2019 | Yes | No |
| Results article | | 09/11/2021 | 26/10/2021 | Yes | No |
| Results article | Under-detection of blood culture-positive enteric fever cases: The impact of missing data and methods for adjusting incidence estimates | 16/01/2020 | 11/10/2023 | Yes | No |
| Results article | Immune responses to typhoid conjugate vaccine in a two dose schedule among Nepalese children <2 years of age | 22/02/2024 | 17/10/2025 | Yes | No |
| Protocol article | protocol | 07/03/2019 | 08/03/2019 | Yes | No |
| Participant information sheet | Participant information sheet | 11/11/2025 | 11/11/2025 | No | Yes |