Premedication for intubation in neonates: a randomised controlled trial

Submission date 19/10/2004	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date	Overall study status	Statistical analysis plan
19/10/2004	Completed	[X] Results
Last Edited 25/08/2009	Condition category Surgery	Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s) Scientific

Contact name Dr Brigitte Lemyre

Contact details

CHEO, Dept of Pediatrics 401, Smyth Road Ottawa Canada K1H 8L1 +1 613 737 8561 blemyre@ottawahospital.on.ca

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers N/A

Study information

Scientific Title

Acronym

PIN

Study objectives

Elective endotracheal intubations are still commonly performed without premedication in many institutions. The hypothesis tested in this study was that morphine given prior to elective intubations in neonates would decrease fluctuations in vital signs, shorten the duration of intubation and reduce the number of attempts.

Ethics approval required

Old ethics approval format

Ethics approval(s) Not provided at time of registration

Study design Randomised controlled trial

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) Hospital

Study type(s) Other

Participant information sheet

Health condition(s) or problem(s) studied Endotracheal intubation in neonates

Interventions Morphine 0.2 mg/kg IV compared to placebo 5 minutes before an endotracheal intubation

Intervention Type Procedure/Surgery

Phase Not Specified

Primary outcome measure

The study aimed to test the hypothesis that morphine 0.2 mg/kg would decrease fluctuations in vital signs, shorten the duration of the procedure and reduce the number of attempts. The primary outcome was the duration of severe hypoxemia, defined as Sp02 < 85% with a HR< 90 /min. This was felt to be the most undesirable side effect of endotracheal intubation as cerebral blood flow in neonates is highly dependent upon heart rate.

Secondary outcome measures

- 1. Duration of the procedure
- 2. Duration of hypoxemia (Sp02 < 85%)
- 3. Number of attempts
- 4. Maximum change in blood pressure from baseline
- 5. Occurrence of bradycardia (HR<90/min).

Overall study start date

01/12/1999

Completion date

30/09/2000

Eligibility

Key inclusion criteria

Newborn infants of all gestations admitted to one Neonatal Intensive Care Unit. Infants of all gestations, admitted to McMaster University Medical Center level III NICU and considered likely to need an elective oral or nasotracheal intubation during their hospital stay, were candidates for inclusion in this study. Families were approached for consent as soon as possible after birth when an elective intubation during their hospital stay seemed likely: if their infant(s) was less than 30 weeks gestation, already ventilated (as endotracheal tubes are frequently changed after 10 days if clinical deterioration from a respiratory standpoint), was on NCPAP for respiratory distress or was needing an elective surgery. Others were approached when an elective intubation was needed. At the time of this study, our unit was a 33-bed level 3 NICU, caring for both inborn and outborn patients, and the referral center for 25000 annual deliveries, with 900-1000 admissions per year.

Participant type(s)

Patient

Age group

Neonate

Sex Both

Target number of participants 34

Key exclusion criteria

- 1. Absence of an intravenous access
- 2. Upper airway anomaly potentially leading to a difficult intubation
- 3. Cyanotic heart disease

4. Upper gastrointestinal obstruction (which would require a rapid sequence intubation)

5. Concurrent opioid administration.

Date of first enrolment 01/12/1999

Date of final enrolment 30/09/2000

Locations

Countries of recruitment Canada

Study participating centre CHEO, Dept of Pediatrics Ottawa Canada K1H 8L1

Sponsor information

Organisation McMaster University (Canada)

Sponsor details Division of Newborn Medicine 1200 Main Street West Hamilton, ON Canada L8S 4J9

Sponsor type University/education

ROR https://ror.org/02fa3aq29

Funder(s)

Funder type Not defined

Funder Name Not provided at time of registration

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	05/10/2004		Yes	No