Randomised controlled trial on the impact of collateral damage and disconnections on memory outcome after temporal lobe epilepsy surgery and the relevance of surgically caused memory decline for every day activities and functioning

Submission date	Recruitment status	Prospectively registered		
30/09/2008	No longer recruiting	☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
31/10/2008	Completed Condition category	ResultsIndividual participant data		
Last Edited				
31/10/2008	Nervous System Diseases	Record updated in last year		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Johannes Schramm

Contact details

Klinik für Neurochirurgie Sigmund-Freud-Str. 25 Bonn Germany 53105

Additional identifiers

Protocol serial number

Sonderforschungsbereich (SFB) TR3 subproject A1

Study information

Scientific Title

Subtemporal versus transsylvian approach for selective amygdala-hippocampectomy: a randomised controlled trial on the epileptological and cognitive outcome

Study objectives

The aim of the study is to evaluate pre- and post-operative verbal episodic memory, frontal lobe executive functions and reward learning as a function of surgically-caused cortical lesions and dissections of fibre tracts from and to the temporal lobe structures.

As the main surgical variation will be that of a preservation or dissection of the temporal stem, using a subtemporal versus a transsylvian approach, differences due to this variation are expected:

- 1. Preserving the temporal stem (subtemporal approach) results in a better crosstalk between frontal executive and temporal memory functions. Therefore memory aspects of organisation, differentiation, recognition (source memory) theoretically can be expected to be better preserved after the subtemporal approach as compared to the transsylvian approach.
- 2. Partial disconnection of the temporal stem (transsylvian approach) is possibly expected to result in a more pronounced effect of reward learning as the importance of the functional connection between temporal and frontal structures for this function has been discussed recently
- 3. The relevance of cognitive development and seizure control for quality of life and their implication for the patients' every day life are another topic of interest. How much memory decline will be accepted in favor of seizure freeness? With an unfavourable outcome in terms of seizure as well as memory outcome one might assume a negative impact on the patients' memory-dependent activities and their quality of life that might go along with higher rates of depression.

Ethics approval required

Old ethics approval format

Ethics approval(s)

University of Bonn Medical Centre Ethics Committee approval received on the 30th April 2008 (ref: 032/08)

Study design

Prospective, interventional, randomised single-centre study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Intractable mesial temporal lobe epilepsy (MTLE)

Interventions

As part of the pre-surgical evaluation patients and healthy controls will be neuropsychologically tested and investigated with functional as well as structural MRI. Patients were then randomised to either a subtemporal or a transsylvian surgical approach, resulting in a preservation (subtemporal approach) or dissection (transsylvian approach) of the temporal stem. Post-operatively, patients are seen for MRI-scanning, neuropsychological testing and medical consultation 3, 6 and 12 months after surgery.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Neuropsychological outcome in memory and non-memory functions as well as in brain activation in three fMRI paradigms will serve as dependent measures. Each patient will undergo comprehensive neuropsychological testing pre-operatively and 12 months post-operatively. The following tests will be applied:

- 1. Verbal learning and memory: two parallel versions of a pre- and post-operative Verbal Learning Test (VLMT- Verbaler Lern- und Merkfähigkeitstest). The VLMT (German adaptation of the Rey Auditory Verbal Learning Test) requires five trials of learning and recall of a word list consisting of 15 words, free recall immediately after distraction (learning/recall of a second list in one trial) and a recall after a half-hour delay, which is followed by a recognition trial (list with original words plus distractors).
- 2. Figural learning and memory was obtained using the DCS-R, a German, revised version of the DCS, a design list learning test (Diagnostikum für Zerebralschädigung)
- 3. Language functions:
- 3.1. "Confrontation naming", Boston Naming Test
- 3.2. "Phonematic" and "semantic fluency"
- 3.3. "Token Test", a subtest of the Aachener Aphasie-Test (a German test battery for aphasia) which is seen to measure verbal comprehension
- 4. Attention functions:
- 4.1. "d2-Test", a letter cancellation test
- 4.2. The c.I.T., a short test to measure cerebral insufficiency (Kurztest für cerebrale Insuffizienz)
- 5. Psychomotor speed, mental tracking and cognitive flexibility:
- 5.1. "Trail Making Test A and B" (TMT-A/B)
- 5.2. Motoric sequences after Lurija
- 5.3. Purdue Pegboard
- 5.4. Finger Tapping Test
- 6. Visual and spatial abilities:
- 6.1. Subtest LPS-7 of the Leistungsprüfsystem (LPS), a German intelligence battery
- 6.2. The Mosaic-Test, a subtest of the German version of the Wechsler Adult Intelligence Scale Revised (HAWIE-R)
- 6.3. Labyrinth test
- 7. FMRI investigation: a new fMRI paradigm will be established for reward learning

All respective measures (primary and secondary outcome measures) will be obtained preoperatively and 12 months post-operatively.

Key secondary outcome(s))

These address the question of the patients' quality of life, personality, depression and memory-dependent activities, which will be assessed using the following questionnaires:

- 1. A German version of the Beck Depression Inventory (BDI)
- 2. FPZ (Fragebogen zur Persönlichkeit bei zerebralen Erkrankungen), an unpublished German CNS-disease related personality questionnaire
- 3. A German adaptation of the QOLIE-10 (Quality of Life Inventory in Epilepsy, 10-Item version), a brief questionnaire to screen for quality of life in epilepsy
- 4. "Memory activities in every day life" MADL, a questionnaire on frequencies of memory dependent everyday activities. The MADL, is a new, unpublished rating scale which has been developed as part of a workshop of the International League against Epilepsy in Venice and is intended to fill the gap between neuropsychological memory impairment, subjective opinion and memory in every day life.

All respective measures (primary and secondary outcome measures) will be obtained preoperatively and 12 months post-operatively.

Completion date

30/06/2012

Eligibility

Key inclusion criteria

- 1. Both males and females
- 2. Patients suffering from intractable temporal lobe epilepsy
- 3. Drug resistance
- 4. Presurgical evaluation led to the recommendation of selective amygdalo-hippocampectomy (SAH)
- 4. Only cases with an magnetic resonance imaging (MRI)-based diagnosis of Ammon's horn sclerosis will be included
- 5. Adults (greater than 18 years)
- 6. Patients have to be able to understand the study plan and give written informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Previous temporal lobe surgery
- 2. Inability to undergo neuropsychological testing because of retardation or insufficient German

language skills

- 3. Gravidity
- 4. Claustrophobia
- 5. Metallic implants/non-removable piercings
- 6. Neurological/psychiatric illness

Date of first enrolment

15/10/2008

Date of final enrolment

30/06/2012

Locations

Countries of recruitment

Germany

Study participating centre Klinik für Neurochirurgie

Bonn Germany 53105

Sponsor information

Organisation

German Research Council (Deutsche Forschungsgemeinschaft [DFG]) (Germany)

ROR

https://ror.org/018mejw64

Funder(s)

Funder type

Research council

Funder Name

German Research Council (Deutsche Forschungsgemeinschaft [DFG]) (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes