A randomised controlled study to assess the emotional needs and efficacy of a service to meet identified emotional needs of individuals with macular disease

It status
ecruiting [] Protocol
dy status 📋 Statistical analysis plan
Results
ategory [] Individual participant data
Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Barbara Hedge

Contact details

Department of Clinical Psychology Torbay Hospital Lawes Bridge Torquay United Kingdom TQ2 7AA

Additional identifiers

Protocol serial number

Version 1, 27th July 2007

Study information

Scientific Title

Study objectives

To compare the emotional wellbeing of individuals newly registered as sight impaired with macular disease who have access to an emotional support package of care together with current standard care, with those who only receive the current standard care. The hypothesis is that the additional care group will experience less psychological pathology (anxiety and depression) and higher quality of life compared to the 'standard care' group.

Primary objective is to assess the efficacy of an emotional support service for individuals who are visually impaired.

Secondary objectives include:

- 1. To assess the emotional needs of individuals with macular disease over the 12 months following registration as sight impaired
- 2. To identify factors that predict individuals who are experiencing emotional difficulties adjusting to registration
- 3. To use the findings of the study to inform an update of the Certificate of Visual Impairment (CVI) form

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval pending from the Devon and Torbay Research Ethics Committee as of 24/09 /2007.

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Age related macular disease

Interventions

All patients (both intervention and control groups) will have current standard care plus an initial assessment via completion of questionnaires and an interview administered via the psychology service.

Each of the participants in the intervention group will be allocated to a volunteer who also has a macular disease. The participants will then receive a telephone call from the volunteers they are assigned to, inviting them to attend a 'newly diagnosed' group meeting run by volunteers with macular disease.

Key activities of the newly diagnosed group:

The group aims to:

- 1. Provide information:
- 1.1. Answer questions about macular disease from volunteers' own experiences
- 1.2. Distribute information (leaflets, audio tapes, etc.,) regarding macular disease, the low vision clinic, support services, benefits, etc.,
- 1.3. Describe, invite to and answer questions about the Macular Disease Support Group
- 2. Screen individuals for evidence of unmet needs:
- 2.1. Those who are in low mood, extremely anxious
- 2.2. Those who need but are not obtaining available social service support
- 2.3. Those not accessing appropriate medical services, e.g., low vision clinic
- 3. Provide a link between newly diagnosed individuals and the professional and voluntary services:
- 3.1. Relay questions that they could not answer to the ophthalmologist, social services, clinical psychology, etc., as relevant and ensure that the answer is fed back to the questioner
- 3.2. Relay details of those requiring additional support to either the ophthalmology service, social services or clinical health psychology service as appropriate

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Psychological pathology, measured at baseline, 4 and 12 months:

- 1. Psychiatric status, assessed by the 28-item General Health Questionnaire (GHQ-28)
- 2. Hospital Anxiety and Depression Scale (HADS)
- 3. Risk of deliberate self-harm, assessed by Beck Hopelessness Scale (BHS)

Key secondary outcome(s))

The following secondary outcomes will be measured at baseline, 4 and 12 months:

- 1. Quality of Life, assessed by the Macular Disease-dependent Quality of Life (MacDQoL) scale
- 2. Adjustment, assessed by the Impact of Events Scale (IES)
- 3. Coping strategies, assessed by the COPE questionnaire
- 4. Social support, assessed by the Social Support Questionnaire (SSQ)
- 5. Knowledge, support and satisfaction questionnaire

Completion date

31/01/2011

Eligibility

Kev inclusion criteria

- 1. Patients who are experiencing sight impairment through Age related Macular Disease (AMD) that requires initial registration as sight impaired
- 2. Voluntary informed consent given

Participant type(s)

Patient

Healthy volunteers allowed

Age group

Adult

Sex

All

Key exclusion criteria

- 1. Limited understanding of English
- 2. Diagnosis of dementia
- 3. Have severe learning disabilities
- 4. Have a current severe psychiatric illness
- 5. Sight impairment for reasons other than macular disease
- 6. Unable to give voluntary informed consent

Date of first enrolment

01/10/2007

Date of final enrolment

31/01/2011

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Department of Clinical Psychology

Torquay United Kingdom TQ2 7AA

Sponsor information

Organisation

South Devon Healthcare NHS Foundation Trust (UK)

ROR

https://ror.org/05374b979

Funder(s)

Funder type Charity

Funder Name

Torbay Medical Research Fund (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration