

# Using annual reminders to facilitate uptake of bowel scope screening in previous non-participants: A Randomised Controlled Trial

<b>Submission date</b> 19/06/2015	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 25/08/2015	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 13/11/2019	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background & aims:

Bowel cancer is the fourth most common cancer in England, and the second leading cause of cancer deaths. Detecting the disease early is associated with improved chances for survival; however, most cases are diagnosed late as most patients don't tend to develop symptoms in the early stages. Screening helps to detect cases early, by testing for the disease before any symptoms occur. In England, there is a national bowel cancer screening programme where men and women are invited for a 'one-off' test, called a flexible sigmoidoscopy (or 'bowel scope'), shortly after their 55th birthday. This test is able to save lives from bowel cancer not only by detecting cases early, by preventing them through the timely removal of growths which pre-cancerous (these are commonly referred to as polyps or 'adenomas'). However, less than half of all people (around 40%) attend their screening appointment when invited by the NHS, and this limits the number cancers the programme is able to detect/prevent. There are a number of reasons why people do not screening appointments, such as forgetting to go, being worried that the screening test might be embarrassing or painful and not being unavailable to attend at that time. Anyone who does not attend their appointment can re-book up until the age of 60, but is otherwise not re-invited to participate (people only ever receive one invitation, it is then up to them to contact the screening centre if they want to arrange an appointment). The aim of this study was to examine whether mailed reminders, prompting non-participants to self-refer for screening one & two years after their invitation, could increase screening rates, and whether these 'annual reminders' are more effective when combined with locally tailored information leaflets that have been designed to address patient-related factors for non-attendance (such as worry about the test being painful or embarrassing).

### Who can participate?

Adults aged 55-59 registered with a GP Practice in the London Boroughs of Brent and Harrow.

### What does the study involve?

In this study, participants are randomly assigned to one of three groups: a 'control group', in which participants do not receive a reminder prompting them to book an appointment; a reminder plus 'standard leaflet' group, in which participants receive a reminder and standard

information leaflet one and (if they still have not booked an appointment) two years after their initial invitation; and a reminder plus locally tailored information leaflet group, in which participants receive a reminder and locally tailored information leaflet designed to address common reasons for non-participation one and (if they still have not booked an appointment) two years after their initial appointment. In addition, participants who attend an appointment are asked to complete a questionnaire about why they did not attend previously and why they decided to make attend an appointment at this time.

What are the possible benefits and risks of participating?

Participants aren't able to enrol in the study, they will be randomly selected from a list of screening eligible 'non-participants'. Participants who made a misinformed (i.e. did not understand the purpose of the screening test) or non-deliberate (i.e. forgot to go to their appointment) decision to not participate in screening previously will benefit from a second opportunity to participate, have their concerns about screening addressed and their risk of getting/dying from bowel cancer being reduced. There are risks associate with flexible sigmoidoscopy screening, including bowel perforation (1 in 3,000), which results in serious bleeding.

Where is the study run from?

St. Marks' Hospital, London (UK)

When is study starting and how long is it expected to run for?

March 2015 to September 2016

How long will the trial be recruiting participants for?

St. Marks' Hospital, London (UK)

Who is the main contact?

Sarah Marshall

sarahmarshall5@nhs.net

## Contact information

**Type(s)**

Scientific

**Contact name**

Mr Robert Kerrison

**Contact details**

1-19 Torrington Place

London

United Kingdom

WC1E 7HB

+44 (0)20 7679 8254

robert.kerrison.13@ucl.ac.uk

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

14/0863

## **Study information**

### **Scientific Title**

Use of a locally tailored information leaflet and mailed reminders to facilitate uptake of flexible sigmoidoscopy (bowel scope) screening in previous non-participants: A three-armed Randomised Controlled Trial.

### **Acronym**

ARfFSS (Annual Reminders for Flexible Sigmoidoscopy Screening)

### **Study objectives**

Receipt of a mailed reminder and locally tailored information leaflet will facilitate uptake in previous flexible sigmoidoscopy (bowel scope) screening non-participants. The reminder will be more effective when sent with a locally tailored information leaflet, rather than the national leaflet. A second reminder delivered one year later will increase uptake further.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Ethics Board: NRES Committee North East - Tyne & Wear South

Reference: 15/NE/0043

Date study approved: 30th January 2015

### **Study design**

A single centre, three-armed, Randomised Controlled Trial.

### **Primary study design**

Interventional

### **Secondary study design**

Randomised controlled trial

### **Study setting(s)**

Hospital

### **Study type(s)**

Screening

### **Participant information sheet**

Not available in web format, please use contact details to request a participant information sheet

## **Health condition(s) or problem(s) studied**

We are investigating whether a mailed reminder, with instructions on how to book an appointment, can facilitate uptake of flexible sigmoidoscopy (bowel scope) screening in previous non-participants

## **Interventions**

Trial Arm 1: Control (i.e. usual care, which is invitation to participate in BSS at 55 only - participants randomly allocated to this arm of the study will not receive anything additional to usual care as a result of this study).

Trial Arm 2: Intervention Arm A (Participants randomly allocated to this group will receive a mailed reminder stating they previously did not attend BSS, but can self-refer up until the age of 59. The mailed reminder will include instructions on how to book an appointment, as well as an 'appointment request slip' which the patient can return to St. Marks' screening centre using the free-post envelope provided. A copy of the national information leaflet will also be provided. The intervention will be delivered a second time one year later, for those participants who have still not attended an appointment).

Trial Arm 3: Intervention Arm B (Participants randomly allocated to this group will receive a mailed reminder stating that they previously did not attend BSS, but can self-refer up until the age of 59. The mailed reminder will include instructions on how to book an appointment, as well as an 'appointment request slip' which the patient can return to St. Marks' screening centre using the free-post envelope provided. Participants in this arm will receive a locally tailored information leaflet, designed specifically to address barriers to BSS, rather than a copy of the national information leaflet. The intervention will be delivered a second time one year later, for those participants who have still not attended an appointment).

In both intervention groups, participants who attend an appointment will be asked to complete a questionnaire about why they previously did not attend BSS, and why they decided to attend on this occasion.

## **Intervention Type**

Behavioural

## **Primary outcome measure**

1. Uptake in each of the trial arms
2. Uptake in each trial arm between genders
3. Uptake after one round of 'annual reminders'
4. Uptake after two rounds of 'annual reminders'

## **Secondary outcome measures**

1. Patient preferences for a same sex practitioner by gender
2. Reasons for previously not participating in BSS
3. Reasons for participating in BSS on this occasion

## **Overall study start date**

16/03/2015

## **Completion date**

25/09/2016

## **Eligibility**

### **Key inclusion criteria**

Adults who meet the study-specific eligibility criteria (listed below) will be identified on the NHS Bowel Cancer Screening Programme's in-house IT system, BCSS (Bowel Cancer Screening System) by a member of the direct patient care team at St. Marks' Hospital:

1. Must be 56-58 years of age at the time they are enrolled in the study
2. Must be registered with a GP practice served by the St Marks BCSC (i.e. a GP practice in the London Boroughs of Brent and Harrow)
3. Must have previously been offered, but not attended, a routine BSS appointment at least one year ago.
4. Must meet the clinical eligibility criteria for BSS (see exclusion criteria, below)

**Participant type(s)**

Other

**Age group**

Adult

**Sex**

Both

**Target number of participants**

1,386

**Total final enrolment**

1218

**Key exclusion criteria**

Exclusion Criteria:

The St Marks BCSC will exclude any individuals who do not meet the clinical-specific criteria for BSS (as listed below) from the study. These include:

1. Individuals who have had their large bowel removed
2. Individuals who have a stoma bag to collect their stool
3. Individuals currently being treated (for example, with steroids) for inflammatory bowel disease (i.e. ulcerative colitis or Crohn's disease)
4. Individuals who are awaiting heart surgery or who have had heart surgery in the last three months

**Date of first enrolment**

16/03/2015

**Date of final enrolment**

25/07/2016

**Locations****Countries of recruitment**

England

United Kingdom

**Study participating centre**  
**St. Marks' Hospital**  
Watford Road  
Harrow  
London  
United Kingdom  
HA1 3UJ

## **Sponsor information**

### **Organisation**

University College London Joint Research Office

### **Sponsor details**

Joint Research Office, Floor 1, Maple House  
149 Tottenham Court Road  
London  
England  
United Kingdom  
W1T 7DN

### **Sponsor type**

University/education

### **Website**

N/A

### **ROR**

<https://ror.org/02jx3x895>

## **Funder(s)**

### **Funder type**

University/education

### **Funder Name**

St. Marks' Hospital, London (UK)

### **Funder Name**

University College London

### **Alternative Name(s)**

University College London in United Kingdom, Collegium Universitatis Londinensis, UCL

### **Funding Body Type**

Government organisation

### **Funding Body Subtype**

Universities (academic only)

### **Location**

United Kingdom

## **Results and Publications**

### **Publication and dissemination plan**

We plan to disseminate the findings of this study by way of academic conferences and publication in peer-reviewed journal articles.

### **Intention to publish date**

31/12/2016

### **Individual participant data (IPD) sharing plan**

### **IPD sharing plan summary**

Not expected to be made available

### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results of feasibility study	29/03/2016		Yes	No
<a href="#">Results article</a>	results of first 14 months	01/06/2016		Yes	No
<a href="#">Results article</a>	overall results	01/01/2017		Yes	No
<a href="#">Results article</a>	results	22/10/2018	13/11/2019	Yes	No
<a href="#">HRA research summary</a>			28/06/2023	No	No