Total disc replacement versus fusion in cervical radiculopathy

Submission date 17/02/2012	Recruitment status No longer recruiting
Registration date 19/03/2012	Overall study status Completed
Last Edited 20/12/2021	Condition category Musculoskeletal Diseases

[] Prospectively registered

[] Protocol

[] Statistical analysis plan

[X] Results

[] Individual participant data

Plain English summary of protocol

Background and study aims

Cervical radiculopathy is where a slipped disc or other bone pinches or irritates a nerve in the cervical spine (neck), resulting in pain. For patients with severe pain that lasts for more than 3 months, surgery can be effective and reduce symptoms. The standard procedure is to decompress the nerve by removing bone or disc material, and fuse the disc joint together. This is an accepted method with good results. Fusion may put more stress on the adjacent discs and may lead to increased degeneration (damage). Artificial discs have been developed as an alternative to fusion and may reduce degeneration and future problems at adjacent discs. The aim of this study is to compare the results of cervical fusion to disc replacement with an artificial disc implant.

Who can participate?

Patients between 18-60 years of age with cervical radiculopathy who have had symptoms for at least 3 months

What does the study involve?

Patients are randomly allocated to undergo decompression with either fusion or disc replacement. All participants are followed up to assess their symptoms after 3 months and 1, 2 and 5 years.

What are the possible benefits and risks of participating?

The use of artificial discs could prevent changes in adjacent discs and therefore lead to a better long term result. As disc replacement is a quite new method there are some potential risks, such as loosening or dislocation of the implant which might lead to further surgery.

Where is the study run from?

- 1. Stockholm Spine Center (Sweden)
- 2. Ryhov Hospital (Sweden)
- 3. Uppsala University Hospital (Sweden)

When is the study starting and how long is it expected to run for? May 2007 to May 2012 Who is funding the study? 1. DePuy Spine, Inc. (Sweden) 2. Johnson & Johnson AB (Sweden)

Who is the main contact? Dr Martin Skeppholm

Contact information

Type(s) Scientific

Contact name Dr Martin Skeppholm

Contact details Stockholm Spine Center Löwenströmska Sjukhuset Upplands Väsby Sweden 19489

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers N/A

Study information

Scientific Title

Total disc replacement versus fusion in cervical radiculopathy: a multicenter randomized controlled trial

Study objectives

Null hypothesis: Disc replacement does not give a better outcome than fusion when cervical radiculopathy is treated with surgery.

Ethics approval required Old ethics approval format

Ethics approval(s) Regional Ethics Committee, Stockholm, 31/03/2007, ref: 2006/1266-31/3 **Study design** Multicenter randomized controlled trial

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Cervical radiculopathy

Interventions

1. Decompression and fusion of 1-2 cervical levels. Fusion is done with bone graft from iliac crest and anterior plate stabilisation

2. Decompression and disc replacement with Discover disc prosthesis (DePuy Spine)

Both groups are followed up at 3 months, 1 year, 2 years and 5 years.

Intervention Type

Other

Phase Not Applicable

Primary outcome measure

Neck Disability Index (NDI) at 2 years follow up after intervention

Secondary outcome measures

- 1. Visual Analaogue Scale (VAS) neck/arm
- 2. EQ-5D
- 3. Dysphagia Short Questionnaire (DSQ)
- 4. Radiographic evaluation
- 5. Sick leave
- 6. Consumption of analgetics

Overall study start date

20/05/2007

Completion date

31/05/2012

Eligibility

Key inclusion criteria

At least 3 months of radiculopathy
 Associated findings on Magnetic resonance imaging (MRI) at 1-2 levels in cervical spine
 18-60 years old

Participant type(s)

Patient

Age group Adult

Lower age limit 18 Years

Upper age limit 60 Years

Sex Both

Target number of participants 150

Key exclusion criteria

1. Previous cervical surgery

- 2. Drug abuse or other obvious reason for bad compliance
- 3. History of Whiplash associated disorder (WAD) or severe cervical trauma
- 4. Rheumatoid arthritis, known malignancy, active infection or other systemic disease
- 5. Pregnancy
- 6. Cervical malformations

Date of first enrolment

20/05/2007

Date of final enrolment

31/05/2012

Locations

Countries of recruitment Sweden

Study participating centre

Stockholm Spine Center Upplands Väsby Sweden 19489

Sponsor information

Organisation DePuy Spine (Sweden)

Sponsor details c/o Johnson & Johnson AB Staffans väg 2 Sollentuna Sweden SE-191 84

Sponsor type

Industry

Website http://www.jjnordic.com/sv-se/Franchise/DePuySpine/Sidor/default.aspx

Funder(s)

Funder type Industry

Funder Name DePuy Spine, Inc (Sweden)

Funder Name Johnson & Johnson AB (Sweden)

Results and Publications

Publication and dissemination plan Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<u>Results article</u>	results	11/01/2019	16/01/2019	Yes	No
<u>Results article</u>	follow up results	01/01/2021	22/01/2021	Yes	No
<u>Results article</u>	10-year results	17/12/2021	20/12/2021	Yes	No