Promotion of dietetic treatment with enhanced salt taste using Salt Chip

Submission date 10/07/2019	Recruitment status No longer recruiting	Prospectively registered
		☐ Protocol
Registration date 24/07/2019	Overall study status Completed	Statistical analysis plan
		Results
Last Edited	Condition category	Individual participant data
24/07/2019	Circulatory System	Record updated in last year

Plain English summary of protocol

Background and study aims

Adequate dietary intake to satisfy the nutrition requirement is particularly crucial for the patients who undergo invasive surgery, such as open heart surgery and large vessel surgery, to maintain the nutritional status, reduce edema, and start cardiac rehabilitation as early as possible during the perioperative period. However, this is often hindered by the physical damages in the mouth, reduction of the pylorus movement, and the lowered motivation by the surgery. In addition, dietary restriction of salt with the intake less than 6.0 g/day, which is reported to be effective as a nutrition therapy for cardiovascular diseases, often leads to further reduction of dietary intake due to the low-salt taste.

We consider that dietetic treatment to increase the food intake and thus improve the nutrition status during the perioperative period is of importance for good recovery of the patients. In this clinical trial, we focus on salt since salt is one of the major contributors to good taste while the intake is strictly restricted for the cardiovascular patients. We used Salt Chip to enhance the salt taste without increasing the salt intake. Salt Chip is attached onto the back of the lower front tooth and deliver salt directly to the tongue, which enables the patients to perceive strong salt taste even with as low as 0.05 g of salt. Dietetic treatment with Salt Chip should increase the eating amount of patients who underwent open heart surgery, which will promote recovery of the activities of daily life.

Who can participate?

Patients who undergo open heart surgery without severe frailty.

What does the study involve?

Participants of Group S are asked to use Salt Chip at every meal during their stay at the hospital except the first meal after the surgery. The participants of control group C do not use Salt Chip. The eating rate at each meal, which is 100% when a subject eats all the food provided and 0% when he/she did not eat any, is calculated from the weight of the food on the plates before and after the meal. Albumin (Alb g/dL), c-reactive protein (CRP mg/dL) and total cholesterol (T-cho mg/dL) are measured before the surgery, one day after the surgery (POD1), 7 days after the surgery (POD7), and when leaving the hospital (ENT).

Body composition is measured using InBody S10 (InBody Japan Inc., Tokyo, Japan), which includes body water balance, hydration percentage (%), total protein (g/dL), total fat (kg), skeletal muscle mass (kg), body cell mass (kg), and basal metabolic rate (kcal/day). The body water balance is the volumetric ratio of the extracellular water to the intracellular water. The hydration percentage is the ratio of the extracellular water amount to the body weight without fat. The measurement is conducted at 7:00 am before breakfast from POD1 to POD7 and at 16: 00 pm before dinner in the preoperative period and from POD8 to POD14.

Progress of rehabilitation is evaluated by treadmills exercise. We assume that the subjects recover the activities of daily living (ADL) when they can conduct treadmills exercise for 20 min.

What are the possible benefits and risks of participating?

Enhanced salt taste using Salt Chip may increase the eating rate, which can be beneficial for fast recovery. Participants may percept too much salt taste, in which case the participants can drop out of the trial immediately.

Where is the study run from?

The study is conducted at Osaki Hospital Tokyo Heart Center.

When is the study starting and how long is it expected to run for? March 2017 to June 2017.

Who is funding the study?

Keio University provides In-Body. LTaste Inc. provides Salt Chip. The remaining incidental costs are paid by Osaki Hospital Tokyo Heart Center.

Who is the main contact?

Dr. Minoru Yoshida

minoru.yoshida.tokyoheart@gmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Minoru Yoshida

ORCID ID

https://orcid.org/0000-0002-6711-0306

Contact details

Osaki Hospital Tokyo Heart Center 5-4-12 Kita-Shinagawa Shinagawa Tokyo Japan 141-0001 +81-3-5789-8100 minoru.yoshida.tokyoheart@gmail.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

0084CR

Study information

Scientific Title

Effect of dietetic treatment with enhanced salt taste in cardiovascular surgery perioperative period

Acronym

SaltChip

Study objectives

- 1. Enhanced salt taste promotes eating rate during the perioperative period.
- 2. Increased eating rate is beneficial in recovery from the surgery.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved, the ethics committee of Osaki Hospital Tokyo Heart Center (5-4-12 Kita-Shinagawa, Shinagawa, Tokyo, 141-0001, Japan; +81-3-5789-8100; +81-3-5789-8101), ref: 0084CR.

Study design

Interventional, nonblinded, parallel trial, single-centre

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Open heart surgery patients

Interventions

The participants in the test group were requested to use Salt Chip®, which can enhance the salt taste without increasing the salt intake amount, at every meal.

The duration of the intervention is from one day before the open heart surgery to ENT.

The eating rate at each meal, which is 100% when a subject eats all the food provided and 0% when he/she did not eat any, was calculated from the weight of the food on the plates before and after the meal. Albumin (Alb g/dL), c-reactive protein (CRP mg/dL) and total cholesterol (T-cho mg/dL) were measured before the surgery, one day after the surgery (POD1), 7 days after the surgery (POD7), and when leaving the hospital (ENT).

Body composition was measured using InBody S10 (InBody Japan Inc., Tokyo, Japan), which included body water balance, hydration percentage (%), total protein (g/dL), total fat (kg), skeletal muscle mass (kg), body cell mass (kg), and basal metabolic rate (kcal/day). The body water balance is the volumetric ratio of the extracellular water to the intracellular water. The hydration percentage is the ratio of the extracellular water amount to the bodyweight without fat. The measurement was conducted at 7:00 am before breakfast from POD1 to POD7 and at 16:00 pm before dinner in the preoperative period and from POD8 to POD14.

Progress of rehabilitation was evaluated by treadmills exercise. We assumed that the subjects recovered the activities of daily living (ADL) when they could conduct treadmills exercise for 20 min.

At approximately 2 weeks the patients who visited the hospital for open-heart surgery during the period were randomised to either the salt chip group or control group. The corresponding period was iterated.

In addition, it was checked if there was any significant differences between the groups in terms of: age, sex, ejection fraction (EF), eating rate before the surgery, Alb, T-cho, body water balance, hydration percentage, total protein, total fat, skeletal muscle mass, body cell mass, and basal metabolic rate (p-value > 0.05).

Intervention Type

Other

Primary outcome(s)

The eating rate of the patients during the postoperative period is measured at each meal.

Key secondary outcome(s))

Progress of rehabilitation was evaluated by treadmills exercise every day. We assumed that the subjects recovered the activities of daily living (ADL) when they could conduct treadmills exercise for 20 min.

Completion date

07/07/2017

Eligibility

Key inclusion criteria

Open heart surgery patients

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Sex

All

Total final enrolment

30

Key exclusion criteria

- 1. Unable to give informed consent and/or assent
- 2. Severe frailty

Date of first enrolment

18/03/2017

Date of final enrolment

19/06/2017

Locations

Countries of recruitment

Japan

Study participating centre Osaki Hospital Tokyo Heart Center

5-4-12 Kita-Shinagawa, Shinagawa, Tokyo, Japan Tokyo Japan

141-0001

Sponsor information

Organisation

Osaki Hospital Tokyo Heart Center

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Osaki Hospital Tokyo Heart Center

Funder Name

LTaste Inc.

Funder Name

Keio University

Alternative Name(s)

, , , Keio

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

Japan

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication.

IPD sharing plan summary

Other

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 No Yes