

# Corneal tattooing for corneal opacities

<b>Submission date</b> 20/10/2017	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 26/10/2017	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 25/11/2020	<b>Condition category</b> Eye Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Corneal (the front part of the eye that covers the pupil and iris) tattooing is one of the options offered to patients with corneal opacities (problems that lead to scarring or clouding of the cornea). However, there are many other types of treatments that have impacted the popularity of corneal tattooing. Various tattooing methods have been used such as: chemical dyeing with gold or platinum chloride, and nonmetallic tattooing with Indian ink, Chinese ink, lamp black, and other organic dyes. The aim of this study is to examine if treatment of corneal opacities by painting them with Rotring Chinese ink.

### Who can participate?

Patients with superficial or deep corneal opacity causing severe disfigurement or those who are blind.

### What does the study involve?

The procedure is carried out in the operating room under sterile conditions by one surgeon (AHA) under topical anesthesia in all patients. Corneal epithelium is not removed. The ink is administered by multiple corneal injections with ink pre-loaded from a sterile cup. The number of injections is determined by the density of the scar and ranges from 4-8 injections. Saline solution is applied to irrigate the corneal surface to wash away excess ink and allow good visualization between injections. Contact lens are then applied and removed after one week.

### What are the possible benefits and risks of participating?

Participants may benefit from an improvement in cosmetic appearance of the eyes. There are no expected risks as the maneuver was tried on rabbits before so there is no risk of dissemination or long term complication on the cornea.

### Where is the study run from?

Sohag University (Egypt)

### When is the study starting and how long is it expected to run for?

June 2016 to June 2017

### Who is funding the study?

Sohag University Hospital (Egypt)

Who is the main contact?  
Dr Engy Mostafa

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Engy Mostafa

**ORCID ID**  
<http://orcid.org/0000-0002-5731-1972>

**Contact details**  
Sohag University Hospital  
Ophthalmology Department  
Sohag  
Egypt  
82525

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**  
1

## Study information

**Scientific Title**  
Outcomes of corneal tattooing by Rotring Painting Ink in disfiguring corneal opacities

**Study objectives**  
The aim of this study is to examine if treatment of corneal opacities by painting them with Rotring Chinese ink.

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**  
Ethical Committee of Faculty of Medicine Sohag University, 25/07/2016

**Study design**  
Prospective interventional non-comparative clinical study

**Primary study design**

Interventional

**Secondary study design**

Non randomised study

**Study setting(s)**

Hospital

**Study type(s)**

Quality of life

**Participant information sheet**

No participant information sheet available.

**Health condition(s) or problem(s) studied**

Total corneal leukomas

**Interventions**

Participants receive conreal tatooing. The procedure is carried out in the operating room under sterile conditions by one surgeon (AHA) under topical anesthesia in all patients. Corneal epithelium is not removed. The ink is administered by multiple transepithelial intrastromal corneal injections using a 30 gauge needle attached to an insulin syringe with ink pre-loaded from a sterile cup. The bevel of the needle is up and administered tangential to the corneal surface to end up in approximately in the mid stroma avoiding accidental perforation of the cornea. The number of injections is determined by the density of the scar and ranged from 4-8 injections. Saline solution is applied to irrigate the corneal surface to wash away excess ink and allow good visualization between injections. Contact lens are then applied and removed after one week.

Postoperatively, moxiflocacin and 1% prednisolone acetate eyedrops are prescribed five times per day for two weeks. NSAID are prescribed twice daily for 3 days. Participants are followed up at one day, one week, one, three and six months. Photographs are taken after one month for comparison. Retreatment is done when needed as in inadequate coloration from the start or fading of the color.

**Intervention Type**

Procedure/Surgery

**Primary outcome measure**

Corneal opacity being tattooed is measured using the slitlamp to judge fading and photographing the eyes at day one, one week, and one month.

**Secondary outcome measures**

Postoperative complications is measured using slitlamp at day one, week one and month one and six months.

**Overall study start date**

01/06/2016

**Completion date**

30/06/2017

## Eligibility

**Key inclusion criteria**

1. No specific age
2. No specific gender
3. Superficial or deep corneal opacity causing severe disfigurement
4. Blind eyes

**Participant type(s)**

Patient

**Age group**

All

**Sex**

Both

**Target number of participants**

50

**Total final enrolment**

53

**Key exclusion criteria**

1. Chronically inflamed eyes
2. Severe corneal calcification or neovascularization
3. Phthisical eyes
4. Anterior

**Date of first enrolment**

01/09/2016

**Date of final enrolment**

30/12/2017

## Locations

**Countries of recruitment**

Egypt

**Study participating centre**

Sohag University

Sohag

Egypt

82525

# Sponsor information

## Organisation

Sohag Univerity Hospital

## Sponsor details

Ophthalmology Department

Sohag University

Faculty of Medicine

Sohag

Egypt

82525

## Sponsor type

Hospital/treatment centre

## ROR

<https://ror.org/02wgx3e98>

# Funder(s)

## Funder type

Hospital/treatment centre

## Funder Name

Sohag University Hospital

# Results and Publications

## Publication and dissemination plan

Planned publication in a high-impact peer reviewed journal. The protocol is available on request.

## Intention to publish date

30/10/2017

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Engy Mohamed Mostafa at [engymostafa@yahoo.com](mailto:engymostafa@yahoo.com).

## IPD sharing plan summary

Available on request

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	25/06/2018	25/11/2020	Yes	No