

Femoral nerve blockade in hip fracture patients

Submission date 30/03/2009	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 05/05/2009	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 05/05/2009	Condition category Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Ola Winso

Contact details
Operationscentrum
Norrlands universitetssjukhus
Umea
Sweden
SE-901 85

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title
Femoral nerve blockade in hip fracture patients: a randomised controlled trial

Study objectives
Femoral nerve blockade reduces pain and the need of opioids and therefore also post-operative delirium and complications in hip fracture patients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethical Committee of the Faculty of Medicine at Umea University approved on the 7th October 2008 (ref: 08-121M)

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Hip fracture

Interventions

Patients are randomised to femoral nerve blockade or the regular use of opioids. Both groups will receive 1 g of paracetamol 4 times/day. The patients in the intervention group will receive a femoral nerve blockade as soon as they arrive at the Orthopaedic Department. If the patients in the intervention group assess pain according to Visual Analogue Scale (VAS) more than 4 then they will be given morphine intravenously (iv) according to the standard protocol (morphine 10 mg/ml, 1 - 5 mg when necessary). Patients in the control group will be given morphine iv according to the standard protocol when they assess pain according to VAS more than 4 but no blockade (morphine 10 mg/ml, 1 - 5 mg when necessary). Post-operative pain treatment will be given according to the standard protocol in both arms of the study. The total follow-up of the trial for both arms will end at the time of discharge.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. Post-operative delirium, assessed three times a day at the Orthopaedic Department
2. Post-operative complications, such as decubital ulcers, infections, thrombosis, heart failure, assessed three times a day at the Orthopaedic Department
3. Pain, assessed three times a day at the Orthopaedic Department

A cognitive test will be assessed in the ambulance pre-hospitally and at 24 +/- 6 hours post-operatively. At days three to five a more thorough assessment will be done including delirium, depression, cognitive status, quality of life and more.

Key secondary outcome(s)

1. Mortality
2. Orthopaedic recovery, recorded at the time of discharge from the hospital
3. EQ-5D
4. Economics

A cognitive test will be assessed in the ambulance pre-hospitally and at 24 +/- 6 hours post-operatively. At days three to five a more thorough assessment will be done including delirium, depression, cognitive status, quality of life and more.

Completion date

31/12/2010

Eligibility

Key inclusion criteria

1. Both males and females, aged 70 years and above
2. All hip fracture patients admitted to the Orthopaedic Department

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Key exclusion criteria

1. Local infection
2. Allergic to local anaesthesia
3. Dying patients
4. Pathologic hip fractures

Date of first enrolment

30/03/2009

Date of final enrolment

31/12/2010

Locations

Countries of recruitment

Sweden

Study participating centre

Operationscentrum

Umea

Sweden

SE-901 85

Sponsor information

Organisation

Umeå University (Sweden)

ROR

<https://ror.org/05kb8h459>

Funder(s)

Funder type

Government

Funder Name

County Council of Västerbotten (Västerbottens läns landsting [VLL]) (Sweden)

Funder Name

Umeå University (Sweden)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration