# Catheter ablation for the cure of atrial fibrillation study

| Submission date               | Recruitment status No longer recruiting | <ul><li>Prospectively registered</li></ul> |  |  |
|-------------------------------|---|--|--|--|
| 07/10/2015                    |   | ☐ Protocol                                 |  |  |
| Registration date             | Overall study status                    | Statistical analysis plan                  |  |  |
| 19/10/2015                    | Completed                               | [X] Results                                |  |  |
| <b>Last Edited</b> 20/12/2016 | Condition category Circulatory System   | [] Individual participant data             |  |  |

#### Plain English summary of protocol

Background and study aims

Atrial fibrillation (AF) is a common heart condition, affecting millions of people worldwide. The heart consists of two upper chambers (atria) and two lower chambers (ventricles). Inside the right atrium, a cluster of cells (sinus node) are responsible for firing electrical signals into the heart muscle causing the heart to beat regularly (sinus rhythm). When a person is suffering from AF, the normal signals from the sinus node do not work properly, causing other parts of the atria to fire chaotically. These uncoordinated signals cause the heart to beat irregularly and often very fast (arrhythmia). A common starting point for treating AF is the use of antiarrhythmic drugs. There are several types of antiarrhythmics, however they generally work on the nervous system to prevent abnormal firing of electrical signals. Although these medications are widely used, some people do not respond to treatment. Catheter ablation is a surgical technique which is used to destroy the areas of the heart which are sending out the irregular signals. In the procedure, a thin flexible tube (catheter) is inserted into the groin and guided up to the heart. The tip of the catheter then either burns (radiofrequency ablation) or freezes (cryoablation) the affected areas. This procedure is often very successful, and can completely cure AF in some cases. The aim of this study is to find out whether using catheter ablation at the same time as antiarrhythmic drugs is better at improving sinus rhythm than antiarrhythmic drugs alone.

#### Who can participate?

Adults with a history of AF who are intolerant to antiarrhythmic drugs or treatment with antiarrhythmic drugs has failed.

#### What does the study involve?

Participants are randomly allocated to one of two groups. The first group receive a combination of antiarrhythmic drugs and a catheter ablation procedure. The second group receive antiarrhythmic drugs only. All participants are asked to wear an ECG recorder (a device which monitors how well the heart is beating and if there are any irregularities) for three months, which send information to the research team every day. Participants in both groups are also have regular ECG scans over a period of 144 months to find out whether their sinus rhythm has improved and to monitor how many participants in each group survive all together.

What are the possible benefits and risks of participating?

A potential benefit is that patients who receive catheter ablation may show an improvement to their sinus rhythm and may have a better chance of survival. Risks of participating include the general risks associated with antiarrhythmic drugs, as well as a low risk of complications in the catheter ablation group from the procedure.

Where is the study run from? Casa di Cura San Michele (lead centre) and 3 other hospitals in Italy

When is the study starting and how long is it expected to run for? February 2002 to May 2015

Who is funding the study? Casa di Cura San Michele (Italy)

Who is the main contact? Dr Emanuele Bertaglia

# Contact information

#### Type(s)

Scientific

#### Contact name

Dr Emanuele Bertaglia

#### **ORCID ID**

https://orcid.org/0000-0002-3878-2904

#### Contact details

via Ca' Rossa, 35 Venice Italy 30173

#### Additional identifiers

#### Protocol serial number

NA

# Study information

#### Scientific Title

Catheter ablation treatment in patients with drug-refractory atrial fibrillation: A prospective, multi-centre, randomized, controlled study

#### Acronym

**CACAF** 

#### Study objectives

Concurrent catheter ablation and antiarrhythmic drugs improve maintenance of sinus rhythm in comparison to antiarrhythmic drugs alone in patients with paroxysmal and persistent atrial fibrillation.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

San Michele Clinic - Maddaloni (CE) Italy, 07/12/2001, ref: PROT. 22001/CE

#### Study design

Prospective multi-centre randomised parallel trial

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Atrial fibrillation

#### Interventions

Participants are randomly allocated into two treatment groups:

Group 1: Participants in the first group are given antiaarhythmic drugs + catheter ablation (pulmonary vein ablation, right isthmus ablation, left isthmus ablation).

Group 2: Participants in the first group are given antiaarhythmic drugs (preferably amiodarone) only.

In both arms, treatment with antiarrhytmic drugs lasted for the whole follow-up. Patients were followed up by means of a trans-telephonic ECG recorder (Sorin Life Watch, Italy), with data being transmitted daily for 3 months, and by means of standard ECG, Holter monitoring at 1, 4, 7, 10, and 13 months. Thereafter, patients were followed up according to local practice, and between 1st June 2014 and 31st May 2015 they underwent an in-office examination or a phone interview, and repeated a 12-lead ECG.

#### Intervention Type

Mixed

#### Primary outcome(s)

Maintenance of sinus rhythm at follow-up, measured with transtelephonic ECG for the first 3 months, with 24 hour Holter monitoring for the first 12 months, and with 12-lead ECG for 144 months.

#### Key secondary outcome(s))

Cumulative survival, measured as whether a patient is alive at 144 months follow-up (+/- 3 months).

# Completion date

# **Eligibility**

#### Key inclusion criteria

- 1. Aged between 18 and 80 years
- 2. History of paroxysmal (1 or more episodes of AF a month in the last 12 months, each lasting more than 60 min but less than 7 days, with all episodes terminating spontaneously) or persistent (2 or more episodes of AF in the last 12 months, each lasting more than 7 days before being terminated medically) atrial fibrillation first diagnosed at least 6 months before enrollment
- 3. Intolerant of antiarrhythmic drugs or in whom 2 or more antiarrhythmic drug regimens had failed

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

- 1. Permanent atrial fibrillation (atrial fibrillation was the sole rhythm for the last 12 months)
- 2. Atrial fibrillation secondary to a transient or correctable abnormality, including electrolyte imbalance, trauma, recent surgery, infection, toxic ingestion, and endocrinopathy
- 3. Persistence of atrial fibrillation episodes triggered by another uniform arrhythmia (i.e. atrial flutter or atrial tachycardia) despite previous supraventricular tachycardia ablation
- 4. Intra-atrial thrombus, tumour, or other abnormality precluding catheter insertion
- 5. Wolff-Parkinson-White syndrome
- 6. Heart failure with NYHA class III or IV or left ventricular ejection fraction <35%
- 7. Unstable angina or acute myocardial infarction within 3 months
- 8. Cardiac revascularization or other cardiac surgery within 6 months or with prior atrial surgery
- 9. Renal failure requiring dialysis, or hepatic failure
- 10. An implanted device (pacemaker or cardioverter-defibrillator)
- 11. Left atrial diameter >60 mm

#### Date of first enrolment

01/02/2002

#### Date of final enrolment

30/06/2003

# **Locations**

# Countries of recruitment

Italy

#### Study participating centre Casa di Cura San Michele

Via Montella, 16 Maddaloni (CE) Italy 81024

# Study participating centre Ospedale Civile

Via XXIX Aprile, 2 Mirano (VE) Italy 30035

#### Study participating centre Ospedale Civile

Via Battitore, 7/9 Ciriè (TO) Italy 10073

# Sponsor information

# Organisation

Casa di Cura San Michele

#### **ROR**

https://ror.org/00z3eck08

# Funder(s)

# Funder type

Hospital/treatment centre

#### Funder Name

Casa di Cura San Michele

# **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Data sharing statement to be made available at a later date

## **Study outputs**

| Output type     | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-----------------|---------|--------------|------------|----------------|-----------------|
| Results article | results | 01/04/2017   |            | Yes            | No              |