

# Simvastatin (20 mg) vs ezetimibe (10 mg) plus simvastatin (20 mg) in subjects with hypercholesterolaemia and coronary heart disease (CHD)

<b>Submission date</b> 06/07/2004	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 13/08/2004	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 25/11/2013	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr Paul Quartey

### Contact details

Schering-Plough Ltd  
Shire Park  
Welwyn Garden City  
United Kingdom  
AL7 1TW

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

P03435

# Study information

## Scientific Title

### Study objectives

Not provided at time of registration

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Not provided at time of registration

### Study design

Randomised controlled trial

### Primary study design

Interventional

### Secondary study design

Randomised controlled trial

### Study setting(s)

Not specified

### Study type(s)

Not Specified

## Participant information sheet

### Health condition(s) or problem(s) studied

Hypercholesterolaemia and CHD

### Interventions

Double-blind study involving 6 weeks treatment with a once daily dose of simvastatin (20 mg) plus either ezetimibe (10 mg) or matching ezetimibe placebo. Blood samples will be collected prior to treatment to use as a baseline and at the end of the 6 week treatment period to determine the effect of the treatments on the lipid profiles. These pre and post treatment blood samples will also be analysed for haematology and clinical chemistry parameters for safety assessment purposes. The objective of the study is to compare the post treatment lipid results (primarily LDL-C) with the baseline values between the two treatment groups. In addition, the usual safety assessments (i.e. adverse events) and details of concomitant medications etc. will be collected during the study.

### Intervention Type

Other

### Phase

Not Specified

**Primary outcome measure**

Not provided at time of registration

**Secondary outcome measures**

Not provided at time of registration

**Overall study start date**

01/04/2000

**Completion date**

31/05/2005

## Eligibility

**Key inclusion criteria**

Male or female subjects aged 18-75 years with screening low-density lipoprotein cholesterol (LDL-C) between 3.3 and 4.9 mmol/l

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Upper age limit**

75 Years

**Sex**

Both

**Target number of participants**

Not provided at time of registration

**Key exclusion criteria**

Not provided at time of registration

**Date of first enrolment**

01/04/2000

**Date of final enrolment**

31/05/2005

## Locations

**Countries of recruitment**

England

United Kingdom

**Study participating centre**

**Schering-Plough Ltd**

Welwyn Garden City

United Kingdom

AL7 1TW

## **Sponsor information**

**Organisation**

Schering-Plough UK Ltd

**Sponsor details**

Schering-Plough House

Shire Park

Welwyn Garden City

United Kingdom

AL7 1TW

**Sponsor type**

Not defined

**ROR**

<https://ror.org/00148fb49>

## **Funder(s)**

**Funder type**

Industry

**Funder Name**

Schering-Plough UK Ltd

## **Results and Publications**

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration