

Reduction of leg wound complications following coronary artery bypass grafting

Submission date 12/09/2003	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 12/09/2003	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 26/10/2009	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr M Zafar

Contact details
Cardiothoracic Theatres
Level 06
Derriford Hospital
Plymouth
United Kingdom
PL6 8DH
mhzafar40@hotmail.com

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0185115619

Study information

Scientific Title

Study objectives

To determine the best method for leg wound closure. The choice of wound closure technique may have important implications for post operative healing in these high risk cases, and also have an impact in other patients at lower risk for post operative morbidity.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Not Specified

Participant information sheet

Health condition(s) or problem(s) studied

Coronary artery bypass grafting (CABG)

Interventions

A prospective randomised controlled trial performed at the Southwest Cardiothoracic Centre. Recruitment of patients will be over a period of 6 months. Patients will be followed from the day of operation to the time of discharge from the hospital and for a period of 6 weeks thereafter. Patients will be randomised to two equal sized groups.

Patients will be randomly assigned to one of two groups:

1. Single layer wound closure over Redivac drains
2. 2-layer closure +/- Redivac drains as indicated clinically.

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome measure

Post operative healing

Secondary outcome measures

Not provided at time of registration

Overall study start date

25/10/2002

Completion date

25/04/2003

Eligibility

Key inclusion criteria

80 Patients undergoing CABG surgery will be enrolled into the study

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

80

Key exclusion criteria

Does not match inclusion criteria

Date of first enrolment

25/10/2002

Date of final enrolment

25/04/2003

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

Cardiothoracic Theatres
Plymouth
United Kingdom
PL6 8DH

Sponsor information

Organisation

Department of Health (UK)

Sponsor details

Richmond House
79 Whitehall
London
United Kingdom
SW1A 2NL

Sponsor type

Government

Website

<http://www.doh.gov.uk>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Plymouth Hospitals NHS Trust (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/12/2005		Yes	No