

A randomised trial to determine the best method for delivering talc for the management of malignant pleural effusions in patients with a good performance status

Submission date 24/05/2012	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 28/05/2012	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 06/12/2019	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<http://www.cancerresearchuk.org/cancer-help/trials/a-trial-looking-at-how-treat-fluid-lung-tapps>

Contact information

Type(s)

Scientific

Contact name

Dr Nick Maskell

Contact details

Respiratory Research Unit
Southmead Hospital
Bristol
United Kingdom
BS10 5NB

-
nick.maskell@nbt.nhs.uk

Type(s)

Scientific

Contact name

Dr Rahul Bhatnagar

Contact details

-
-
United Kingdom
-

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

HTA 10/50/42, 2843, UKCRN:12537

Study information

Scientific Title

Evaluating the efficacy of Thoracoscopy And talc Poudrage versus Pleurodesis using talc Slurry:
A randomised, open-label trial to determine the most effective method for the management of malignant pleural effusions in patients with a good performance status

Acronym

TAPPS

Study objectives

Primary research question:

Does thoracoscopy and talc poudrage increase the proportion of patients with successful pleurodesis at three months post-procedure, when compared to standard therapy with chest drain insertion and talc slurry instillation?

Secondary research questions:

1. Does thoracoscopy and talc poudrage reduce the time to pleurodesis failure, measured at three and six months post-procedure, when compared to standard therapy with chest drain insertion and talc slurry instillation?
2. Does fluid drainage and talc poudrage at thoracoscopy improve chest x-ray appearances at 24 hours and at 3 months post-procedure, when compared to standard fluid drainage via chest tube alone?
3. Does thoracoscopy and talc poudrage cause less breathlessness and thoracic pain for the first five days post-procedure, when compared to standard therapy with chest drain insertion and talc slurry instillation?
4. Does thoracoscopy and talc poudrage improve health-related quality of life over the six months post-procedure, when compared to standard therapy with chest drain insertion and talc slurry instillation?
5. Is thoracoscopy and talc poudrage cost effective over six months, when compared to standard therapy with chest drain insertion and talc slurry instillation?

6. Does thoracoscopy and talc poudrage reduce healthcare utilisation during the six months post-procedure, when compared to standard therapy with chest drain insertion and talc slurry instillation?

On 07/01/2015 the overall trial end date was changed from 15/01/2015 to 31/01/2017.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES Committee North West – Preston, 26/06/2012, ref: 12/NW/0467

Study design

Randomised open-label multi-centre trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Malignant pleural effusion

Interventions

1. Small-bore chest drain insertion followed by 4 g graded sterile talc slurry pleurodesis
2. Medical (local anaesthetic) thoracoscopy followed by 4 g graded sterile talc poudrage

Intervention Type

Procedure/Surgery

Primary outcome measure

Number of patients who experience pleurodesis failure up to three months (90 days) post randomisation

Secondary outcome measures

1. Requirement for further pleural procedures up to 6 months post-randomisation, as assessed by two independent, blinded adjudicators. The adjudicator will be provided with relevant radiological images and information regarding the patient's health status, including performance status and Visual Analogue Scores (VAS) scores for breathlessness and thoracic pain
2. Percentage radiographic (chest x-ray) pleural opacification, measured by visual estimation in a

blinded fashion, on the side of the pleurodesis attempt at 24 hours post poudrage or slurry instillation, and at 3 and 6 months post randomisation

3. Self-reported health-related quality of life, as measured using the SF-36 and EQ-5D questionnaires measured at 1 month, 3 months and 6 months post randomisation
4. Self-reported thoracic pain, as measured using VAS scores recorded daily for the first 7 days post randomisation, and then weekly for the duration of trial follow-up
5. Self-reported breathlessness, as measured using VAS scores recorded daily for the first 7 days post randomisation, and then weekly for the duration of trial follow-up.
6. The number of patients with pleurodesis failure up to one month (30 days) post randomisation
7. The number of patients with pleurodesis failure up to six months (180 days) post randomisation
8. All-cause mortality up to six months (180 days) post-randomisation
9. Time to pleurodesis failure, censored at six months (180 days) post randomisation
10. Time from randomisation to hospital discharge
11. Number of days spent as a hospital inpatient up to three months
12. Healthcare resource usage and costs at six months (180 days) post randomisation
13. The costs of performing talc pleurodesis under the two interventions under study
14. Follow-up costs

Overall study start date

15/07/2012

Completion date

31/10/2018

Eligibility

Key inclusion criteria

1. Clinically confident diagnosis of malignant pleural effusion requiring pleurodesis, defined as:
 - 1.1. Pleural effusion with histocytologically proven pleural malignancy OR
 - 1.2. Pleural effusion in the context of histocytologically proven malignancy elsewhere, without a clear alternative cause for fluid OR
 - 1.3. Pleural effusion with typical features of malignancy with pleural involvement on cross-sectional imaging (CT/MRI)
2. Fit enough to undergo local anaesthetic thoracoscopy, as per British Thoracic Society (BTS) guidelines
3. Expected survival >3 months
4. Written, informed consent to trial participation

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

330

Total final enrolment

330

Key exclusion criteria

1. Patients in whom thoracoscopy is the only reasonable approach to making a diagnosis, and in whom such a diagnosis would significantly influence further management
2. Age < 18 years
3. Females who are pregnant or lactating
4. Evidence of extensive lung entrapment on chest X-ray (CXR) or CT, or significant fluid loculation on ultrasound scan, to a level which would normally be a contraindication to attempted talc pleurodesis
5. Insufficient volume or position of pleural fluid on lateral decubitus thoracic ultrasound to safely perform local anaesthetic thoracoscopy without further intervention being necessary
6. Previously documented adverse reaction to talc
7. Clear contraindication to thoracoscopy or chest tube insertion

Date of first enrolment

01/08/2012

Date of final enrolment

24/10/2017

Locations**Countries of recruitment**

England

Scotland

United Kingdom

Study participating centre**Southmead Hospital**

Monks Park Avenue

Bristol

United Kingdom

BS10 5NB

Study participating centre**Nottingham City Hospital**

Hucknall Road

Nottingham

United Kingdom

NG5 1PB

Study participating centre
Musgrove Park Hospital
Taunton
United Kingdom
TA1 5DA

Study participating centre
Churchill Hospital
Headington
Oxford
United Kingdom
OX3 7LE

Study participating centre
Medway Maritime Hospital
Gillingham
United Kingdom
ME7 5NY

Study participating centre
King's Mill Hospital
Mansfield Road
Sutton in Ashfield
Nottingham
United Kingdom
NG17 4JL

Study participating centre
Lancashire Teaching Hospitals NHS Foundation Trust
Fulwood
Preston
United Kingdom
PR2 9HT

Study participating centre
Wythenshawe Hospital
Southmoor Road
Wythenshawe
Manchester
United Kingdom
M23 9LT

Study participating centre
Addenbrooke's Hospital
Hills Road
Cambridge
United Kingdom
CB2 0QQ

Study participating centre
St Thomas' Hospital
Westminster Bridge Road
London
United Kingdom
SE1 7EH

Study participating centre
Doncaster Royal Infirmary
Armthorpe Road
Doncaster
United Kingdom
DN2 5LT

Study participating centre
University Hospital of North Tees
Hardwick
Stockton
United Kingdom
TS19 8PE

Study participating centre
Aintree University Hospital
Liverpool
United Kingdom
L9 7AL

Study participating centre
Southern General Hospital
1345 Govan Road
Glasgow

United Kingdom
G51 4TF

Study participating centre

Milton Keynes Hospital NHS Foundation Trust
Standing Way
Eaglestone
Milton Keynes
United Kingdom
MK6 5LD

Study participating centre

Queen Elizabeth Hospital
Birmingham
United Kingdom
B15 2TH

Sponsor information

Organisation

North Bristol NHS Trust (UK)

Sponsor details

c/o Ms Helen Lewis
Research and Innovation
Floor 3, Learning and Research Building
Southmead Hospital
Bristol
United Kingdom
BS105NB

Sponsor type

Not defined

ROR

<https://ror.org/036x6gt55>

Funder(s)

Funder type

Government

Funder Name

NIHR Health Technology Assessment (HTA) (UK) (ref. 10/50/42)

Results and Publications

Publication and dissemination plan

The protocol will be published in an open access journal. The full trial results will be published in peer reviewed journals and presented at national and international conferences. Trial results will also be disseminated to appropriate patient groups/charities upon completion.

Intention to publish date**Individual participant data (IPD) sharing plan**

The trial was funded by the NIHR HTA programme, who will publish the full data and a comprehensive study report at the same time as the academic manuscript. This will be open access and thus available to anyone in perpetuity. For further information email the study Chief Investigator, Nick Maskell, at nick.maskell@bristol.ac.uk. Consent was obtained and all data are anonymised.

IPD sharing plan summary

Other

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Other publications	strategies	21/11/2014		Yes	No
Protocol article	protocol	26/11/2014		Yes	No
Results article	results	05/12/2019	06/12/2019	Yes	No