A feasibility study of electron radiotherapy delivered during a surgical operation for rectal cancer

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
13/05/2022		☐ Protocol		
Registration date	Overall study status Completed Condition category Cancer	Statistical analysis plan		
13/05/2022		Results		
Last Edited		Individual participant data		
09/04/2025		[X] Record updated in last year		

Plain English summary of protocol

Background and study aims

Rectal cancer is a cancer that occurs in the pelvis from the rectum. Locally advanced rectal cancer outgrows the rectum and attaches to other body parts in the pelvis and locally recurrent rectal cancer is a rectal cancer that comes back after surgery, and usually attaches to many different pelvic structures. They are both difficult to manage. The standard of care treatment involves chemotherapy and radiotherapy, followed by what is known as an extended margin operation to remove all cancer affected organs and not leave any cancer cells behind. If cancer cells reach the edge of the removed tissue, there is a high chance of leaving cancer cells behind. This is a key predicter of negative outcome in patients. Intraoperative electron beam radiotherapy (IOERT) was developed to help improve patient outcomes. Once the cancer has been removed, the surgeon and a cancer radiotherapy specialist examine the patient's scans, the cancer specimen and the area the cancer was in, and if there is concern about small numbers of cancer cells being left behind they treat the area with radiotherapy to destroy these cells.

Who can participate?

Patients aged 16 years or older, that are due to receive treatment for the above subsets of rectal cancer will be approached to take part.

What does the study involve?

If eligible on the day of surgery the patient will be randomised to one of three arms: Arm A – standard of care (No IOERT), Arm B – extended margin surgery plus IOERT (10 Gy), or Arm C – extended margin surgery plus higher dose IOERT (15 Gy). The surgeon, cancer specialist team and patient will be blinded to study treatment. Patients will be followed up at 30 days, 3 months and for a minimum of 12 months post surgery as part of the trial and they will be followed up for 5 years as part of standard care.

What are the possible benefits and risks of participating?

The information that we get from this study may help us to treat future patients with the same condition in a more effective way. You will also be helping to further our knowledge of how to treat cancer and this will also benefit society as a whole.

As with every treatment plan, there are associated risks. The risk from surgery includes bleeding, wound infections and a rare chance of death which can occur in less than 1% of people who require surgery. However, all patients will have consented to surgery which is the standard of care. The associated risk from the intraoperative radiotherapy is low and includes wound infections, pain from pelvic nerves, and a narrowing of the tube which carries urine from the kidney to the bladder. The risks of IOERT are counterbalanced however by the potential benefits of better cancer treatment and less chance of cancer recurrence and thereby potential survival.

Where is the study run from? University Hospital Southampton NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for? April 2021 to April 2025

Who is funding the study?
IntraOp Medical Corporation (USA)
PLANETS cancer charity (UK)

Who is the main contact?
Daniel Griffiths, Dpg1e20@soton.ac.uk

Contact information

Type(s)

Public

Contact name

Mr Daniel Griffiths

Contact details

MP131, Southampton General Hospital Tremona Road Southampton United Kingdom SO16 6YD +44 23 8120 5154 electra@soton.ac.uk

Type(s)

Scientific

Contact name

Prof Alex Mirnezami

ORCID ID

https://orcid.org/0000-0002-6199-8332

Contact details

Southampton General Hospital, Mailpoint 824 Somers Cancer Research Building Tremona Road Southampton United Kingdom SO16 6YD

_

A.H.Mirnezami@soton.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

292873

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CPMS 52098, IRAS 292873

Study information

Scientific Title

A double-blinded feasibility study of intraoperative electron radiotherapy in patients with locally advanced or locally recurrent rectal cancer

Acronym

ELECTRA

Study objectives

To determine acceptability and feasibility of recruiting, randomising and delivering IOERT in a randomised controlled trial (RCT) setting as preparatory work for a future late phase RCT.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 31/03/2022, West Midlands - Black Country Research Ethics Committee (The Old Chapel, Royal Standard Place, Nottingham, NG1 6FS, United Kingdom; +44 207 104 8010; blackcountry.rec@hra.nhs.uk), ref: 22/WM/0048

Study design

Interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Colorectal Cancer

Interventions

As part of standard of care, patients with locally advanced or locally recurrent rectal cancer will have an extended margin operation to remove all cancer affected organs.

Participants who meet the eligibility criteria and whom provide their written informed consent for the study will be randomised to one of three study arms: Arm A - extended margin surgery alone (standard of care treatment), Arm B - extended margin surgery with IOERT standard dose (10 Gy), Arm C - extended margin surgery plus IOERT higher dose (15 Gy). Participants will have equal chances to receive one of the three arms of treatment. Randomisation will occur on the day of surgery and the surgeon, cancer specialist team and the patient will be blinded to the result of the randomisation.

The IOERT session will take place in a purposefully designed operating theatre with a portable self-shielded electron beam linear accelerator, known as the MOBETRON. The treating radiation oncologist will determine how the IOERT will be administered based on best practice.

The tests included in the study are a cardiopulmonary exercise test and MRI scans, which are standard of care. A pregnancy test will also be conducted for women of childbearing potential at screening and pre-op, which is not standard of care.

The patients will be asked to complete quality of life and health economics questionnaires at their baseline visit and at 3 and 12 months post surgery.

Participants will be followed up for a minimum of a year in the trial and at the end of the trial they will return to standard care to be followed up until 5 years post surgery. The participants will be followed up at 30 days, 3, 12 and 24 months post surgery as part of the trial.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Feasibility measured at the end of the trial using:

- 1. Number and percentage of patients meeting eligibility criteria and number of patients referred to a specialist MDT over the trial period;
- 2. Number and percentage of patients accepting randomisation;
- 3. Number and percentage of patients for which IOERT was successfully delivered as planned;
- 4. Number and percentage of patients and clinicians for which blinding was maintained for IOERT delivery;
- 5. Percentage of patients whose questionnaires can be analysed;
- 6. Percentage of patients whom we can collect information on potential primary outcomes;

Key secondary outcome(s))

- 1. Morbidity measured using Clavien Dindo at surgery and at 30 days post randomisation
- 2. Mortality at 30 days post randomisation
- 3. Oncological outcomes: IOERT field occurrence, overall recurrence, overall survival at minimum 12 months post randomisation
- 4. Treatment related toxicity graded by CTCAE at minimum 12 months post randomisation
- 5. Time to local or systemic recurrence
- 6. R1 rate at surgery
- 7. Quality of life measured using questionnaires (EQ-5D-5L, LRRC-QoL, SF-36, EORTC QLQ-C30) at pre-op, 3 months and 12 months post surgery
- 8. Resource use and cost at pre-op, 3 months and 12 months post surgery

Completion date

25/04/2025

Eligibility

Key inclusion criteria

- 1. Age >=16 years
- 2. Non-metastatic/oligo-metastatic (up to three lesions from 2 sites predicted to be radically treatable) locally advanced or locally recurrent disease involving the posterior or lateral compartment of the pelvis and predicted to be resectable but with close margins from MRI as determined by a specialist MDT (sMDT)
- 3. Specialist colorectal MDT review with experience in pelvic exenteration, which
- 4. has proposed IOERT as an option for treatment
- 5. Patient suitable for IOERT as a component of treatment in view of the
- 6. responsible Clinical Oncologist
- 7. Performance status <=1 as defined by the Eastern Cooperative Group (ECOG)
- 8. Deemed medically fit for surgery
- 9. Written informed consent.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

16 years

Sex

All

Total final enrolment

31

Key exclusion criteria

Current participant exclusion criteria as of 11/08/2023:

- 1. Unresectable disease/likelihood of R2 resection
- 2. sMDT determined excess prior radiotherapy within IOERT target zone
- 3. Women who are pregnant or breastfeeding
- 4. Participation within an interventional trial within 3 months of the point of registration within ELECTRA

Previous participant exclusion criteria:

- 1. Unresectable disease/likelihood of R2 resection
- 2. sMDT determined excess prior radiotherapy within IOERT target zone
- 3. Women who are pregnant or breastfeeding
- 4. Use of experimental drugs within other interventional clinical trials

Date of first enrolment

20/05/2022

Date of final enrolment

25/04/2024

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Southampton General Hospital

University of Southampton and University Hospital Southampton NHS Foundation Trust Tremona Road Southampton United Kingdom SO16 6YD

Sponsor information

Organisation

University Hospital Southampton NHS Foundation Trust

ROR

https://ror.org/0485axj58

Funder(s)

Funder type

Industry

Funder Name

IntraOp Medical Corporation

Funder Name

PLANETS cancer charity

Results and Publications

Individual participant data (IPD) sharing plan

Individual participant data will be made available, including data dictionaries, for approved data sharing requests. Individual participant data will be shared that underlie the results after deidentification and normalisation of information (text, tables, figures, and appendices). The study protocol and statistical analysis plan will also be available. Anonymous data will be available for request from 3 months after the publication of the results to researchers who provide a completed data sharing request form that describes a methodologically sound proposal, for the purpose of the approved proposal and if appropriate, signed a data-sharing agreement. Pseudonymised participant data within the clinical trial dataset will be available for sharing via controlled access by authorised Southampton Clinical Trials Unit (SCTU) staff. The request for data access will need to detail the specific requirements and the proposed research, statistical analysis, publication plan and evidence of research group qualifications. Data access requests will be reviewed against specific eligibility criteria by the SCTU data custodian and key members of the trial team. Data will be shared once all parties have signed relevant data sharing documentation covering SCTU conditions for sharing and if required, an additional data sharing agreement from the sponsor. Proposals should be directed to ctu@soton.ac.uk.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes