Is it possible to have a chronic obstructive pulmonary disease (COPD) specific breathlessness service in the Netherlands?

Submission date	Recruitment status No longer recruiting	Prospectively registered	
01/11/2021		[X] Protocol	
Registration date	Overall study status	Statistical analysis plan	
06/11/2021	Completed Condition category	Results	
Last Edited		Individual participant data	
05/11/2021	Respiratory	Record updated in last year	

Plain English summary of protocol

Background and study aims

Chronic obstructive pulmonary disease, or COPD, refers to a group of diseases that cause airflow blockage and breathing-related problems. Refractory (treatment resistant) breathlessness is a devastating symptom in COPD. Breathlessness services (BS), usually collaborations of palliative and respiratory teams, offer a multidisciplinary approach. In the Netherlands, few palliative care teams work in an outpatient setting, and BS do not exist.

The study involves training for people with COPD (chronic obstructive lung disease). The aim is to develop training for patients who are troubled by breathlessness. The training cannot take away the breathlessness but should help patients cope with the symptom.

Who can participate?

Patients with COPD and refractory breathlessness.

What does the study involve?

Participants attended at least two sessions with a pulmonologist and a respiratory nurse and one session with a physiotherapist who practiced the breathing techniques with them. The toolkit that was given to patients consisted of a booklet with breathing exercises and a handheld fan. There was not a set number of sessions.

What are the possible benefits and risks of participating?

Benefits: patients who participate might improve their coping skills regarding breathlessness. They also help us set up training that is appropriate for patients. There are no risks in participating; however, the intervention takes up some of the participants' time

Where is the study run from? Spaarne Gasthuis (Netherlands)

When is the study starting and how long is it expected to run for? April 2019 to November 2020

Who is funding the study? Spaarne Gasthuis (Netherlands)

Who is the main contact? Kris Mooren, k.mooren@spaarnegasthuis.nl

Contact information

Type(s)

Scientific

Contact name

Mrs Kris Mooren

ORCID ID

http://orcid.org/0000-0001-7815-0488

Contact details

Spaarne Gasthuis Boerhaavelaan 22 Haarlem Netherlands 2035 RC +31232245969 k.mooren@spaarnegasthuis.nl

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Nil known

Study information

Scientific Title

Testing the waters for a COPD-specific breathlessness service in the Netherlands: a feasibility study

Study objectives

Our primary aim was to demonstrate the feasibility of setting up a BS specifically for COPD patients, with the view of undertaking a randomized controlled clinical trial to test the effectiveness of this BS on breathlessness mastery.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 09/08/2019, Medical Ethics Review Committee of the Amsterdam University Medical Centre (Medisch Ethische Toetsingscommissie VUmc, Van der Boechorststraat 7, kamer H-443, Postbus 7057

1007 MB Amsterdam, Netherlands; +31 20 444 5585; no email provided), ref: 2019.199

Study design

Non-randomized single-center feasibility study

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Quality of life

Participant information sheet

No participant information sheet available

Health condition(s) or problem(s) studied

Improvement of quality of life in COPD patients with refractory dyspnea

Interventions

For this study, the authors were trained by the Cambridge Breathlessness Intervention Service (CBIS). Their Breathing Thinking Functioning model (BTF) was translated into a Dutch toolkit (Ademen Denken Doen, ADD) with full consent of, and in collaboration with, the Cambridge team. The content of the Dutch intervention is similar to the original. However, the service in Cambridge is delivered by occupational therapists, physiotherapists and a consultant in palliative medicine, whereas the Dutch service is delivered by a pulmonologist, a respiratory nurse and a physiotherapist.

The model helps the patients to understand the vicious circles that influence breathlessness. A well-known example is the dyspnea-anxiety-dyspnea cycle: patients have catastrophic thoughts (I need oxygen, I am going to suffocate) (thinking domain), leading to shallow, exhausting 'dead space' breathing (breathing domain), leading to worsening of the symptom that can spiral into a panic attack. These episodes of breathlessness can lead to avoidance of exercise (functioning domain), in turn causing deconditioning that worsens the symptom.

ADD was set up as a brief intervention: patients had at least two sessions with a pulmonologist and a respiratory nurse, and one session with a physiotherapist who practiced the breathing techniques with them. The toolkit that was given to patients consisted of a booklet with breathing exercises and a hand-held fan. If necessary, extra sessions were scheduled. Since we had no experience with giving the intervention in this setting and specifically for COPD patients,

we decided not to fix the number of sessions. However, the average number of sessions for a future randomized multicenter controlled clinical trial shall be derived from this feasibility study.

Intervention Type

Behavioural

Primary outcome measure

Feasibility outcome: number of patients completing the intervention measured using patient records at the end of the study

Secondary outcome measures

- 1. Respiratory problems measured using a Dutch translation of the Chronic Respiratory Questionnaire (CRQ), subset mastery domain at baseline and final visit
- 2. Pulmonary function tests were taken from the patient's medical file at baseline
- 3. Patient and professional satisfaction measured using a postal survey with multiple choice questions after the final visit

Overall study start date

01/04/2019

Completion date

01/11/2020

Eligibility

Key inclusion criteria

Patients with COPD (diagnosed by a pulmonologist, post-bronchodilatator FEV1/FVC below the lower limit of normal) were eligible for the study if they experienced refractory dyspnea (troubled by breathlessness despite optimization of COPD treatment) and were able to visit the outpatient clinic.

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

20

Total final enrolment

19

Key exclusion criteria

- 1. Terminal phase
- 2. Cognitive impairment
- 3. Unable to speak Dutch

Date of first enrolment 01/10/2019

Date of final enrolment 01/11/2020

Locations

Countries of recruitment

Netherlands

Study participating centre Spaarne Gasthuis Boerhaavelaan 22 Haarlem Netherlands

Sponsor information

Organisation

2035RC

Spaarne Gasthuis

Sponsor details

Boerhaavelaan 22 Haarlem Netherlands 2035 RC +31 232240000 wetenschapsbureau@spaarnegasthuis.nl

Sponsor type

Hospital/treatment centre

Website

https://spaarnegasthuis.nl

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Spaarne Gasthuis

Results and Publications

Publication and dissemination plan

Planned publication in a peer-reviewed COPD journal.

Intention to publish date

01/12/2021

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request (k.mooren@spaarnegasthuis.nl)

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol file	version 4	01/08/2019	05/11/2021	No	No