# Perioperative Duloxetine and Etoricoxib improve postoperative pain

Submission date 17/06/2017	<b>Recruitment status</b> No longer recruiting	<ul> <li>Prospectively registered</li> <li>Protocol</li> </ul>
Registration date 19/06/2017	<b>Overall study status</b> Completed	<ul> <li>[_] Statistical analysis plan</li> <li>[X] Results</li> </ul>
Last Edited 26/11/2020	<b>Condition category</b> Surgery	Individual participant data

#### Plain English summary of protocol

Background and study aims

Having spinal surgery requires a lot of recovery time. Usually patients who undergo surgery need medication to help them recover and control any after surgery pain. There are a variety of medications that can be used to help treat patients. Some suggestions for medications to use include duloxetine (which is used to treat anxiety, depression, neurological pain) as well as etoricoxib (an anti-inflammatory (swelling) painkiller). It can be sometimes difficult for healthcare workers to figure out what the best combination of medications is most effective and provides patients with the least amount of side effects while still improving their pain symptoms. The aim of this study is to find the best regimen of drugs for the postoperative (after surgery) analgesia (pain prevention) for patients undergoing spinal surgeries in order to improve patient satisfaction of their postoperative care and to provide them with less side effects from medication after the surgery.

Who can participate?

Adults patients aged 18 and older who are having spinal surgery.

#### What does the study involve?

Participants are randomly allocated to one of four groups. Those in the first group receive a capsule containing a placebo (dummy drug) and two placebo tablets an hour before surgery and then again 24 hours later. Those in the second group receive a capsule a placebo and two etoricoxib tablets an hour before surgery and then again 24 hours later. Those in the first group receive a capsule containing duloxetine and two placebo tablets an hour before surgery and then again 24 hours later. Those in the fourth group receive a capsule containing duloxetine and two placebo tablets an hour before surgery and then again 24 hours later. Those in the fourth group receive a capsule containing duloxetine and two etoricoxib tablets an hour before surgery and then again 24 hours later. Patients then have their surgery as per standard practice. Half an hour after their anaesthesia has worn off and then two, four, six, 12, 24 and 48 hours later, participants are asked to rate their pain levels. Their satisfaction with the treatment they received and pain killer use is also recorded.

What are the possible benefits and risks of participating?

Participants may benefit from being more satisfied with their surgery and less side effects from drugs after the surgery. There are no notable risks with participating however there are risks with any medication and this should be monitored.

Where is the study run from? Minia University Hospital (Egypt)

When is the study starting and how long is it expected to run for? April 2015 to May 2017

Who is funding the study? Investigator initiated and funded (Egypt)

Who is the main contact? Mr Josef Zekry Attia

## **Contact information**

**Type(s)** Scientific

**Contact name** Mr Josef Zekry Attia

**ORCID ID** http://orcid.org/0000-0003-4842-231X

**Contact details** Faculty of Medicine Al-Minia University Main Road Shalaby Land - Menia Al-Minia Egypt 61111 +20 10 00 427407 josefzekry2@yahoo.com

# Additional identifiers

EudraCT/CTIS number

**IRAS number** 

ClinicalTrials.gov number

Secondary identifying numbers 7

# Study information

#### Scientific Title

Perioperative Duloxetine and Etoricoxib to improve postoperative pain after lumbar laminectomy: A randomized, double-blind, controlled study

#### **Study objectives**

Perioperative Duloxetine and Etoricoxib given together are more effective than when given individually at reducing postoperative pain after lumbar laminectomy.

**Ethics approval required** Old ethics approval format

**Ethics approval(s)** Institutional Ethics Committee at Faculty of Medicine Minia university, 01/05/2015, ref: 1-5-2015

**Study design** Four-arm randomised controlled trial

**Primary study design** Interventional

**Secondary study design** Randomised controlled trial

**Study setting(s)** Hospital

**Study type(s)** Treatment

**Participant information sheet** No participant information sheet available

#### Health condition(s) or problem(s) studied

Postoperative pain after laminectomy surguries

#### Interventions

Patients admitted to the hospital are randomised using computer-generated random numbers with closed-sealed envelopes into one of the four groups of 30 patients.

Group 1: Participants receive a placebo capsule + two placebo tablets one hour before surgery and again 24 hours later

Group 2: Participants receive a placebo capsule + 90 mg etoricoxib tablets one hour before surgery and again 24 hours later

Group 3: Participants receive a 60 mg duloxetine capsule + two placebo tablets one hour before surgery and again 24 hours later

Group 4: Participants receive a 60 mg duloxetine capsule + two 90 mg etoricoxib tablets one hour before surgery and again 24 hours later

Participants in all groups undergo surgery as per standard practice.

Participants in all groups are asked to rate their pain levels 30 minutes after the end of anesthesia and two, four, six, 12, 24 and 48 hours post-operatively. In addition, patient satisfaction and pain killer use is also assessed 24 and 48 hours after surgery.

#### Intervention Type

Other

#### Primary outcome measure

Pain is measured using pain assessment measures and questions 30 minutes after the end of anesthesia and two, four, six, 12, 24 and 48 hours post-operatively

#### Secondary outcome measures

1. The time to first rescue analgesic is routinely 24h and 48h postoperatively

2. Total morphine consumption is measured by calculating the doses at 24 and 48 hours postoperatively

3. Side effects, including headache, rash, nausea, vomiting, dizziness and drowsiness are recorded. The severity of postoperative nausea and vomiting (PONV) is graded on a four-point ordinal scale

4. Patient satisfaction is measured using a numerical rating scale (NRS) at 24 hours postoperatively

#### Overall study start date

01/04/2015

### Completion date

01/05/2017

# Eligibility

#### Key inclusion criteria

1. Patients with an ASA physical status of I, II and III scheduled for single level lumbar spinal disc prolapsed surgery

2. Can provide written informed consent

3. Aged between 18 and older

**Participant type(s)** Patient

**Age group** Adult

**Lower age limit** 18 Years

**Sex** Both

#### Target number of participants

131 consecutive patients who met the inclusion criteria were allocated for the study. Eleven patients refused to participate. Therefore, 120 patients were randomized and included in the study. Characteristics of patients and surgical procedures for each group

#### Total final enrolment

120

#### Key exclusion criteria

- 1. History of allergic reaction to any of the study drugs
- 2. History of drug or alcohol abuse
- 3. Abnormal renal or liver function tests.
- 4. Patients using antidepressants who did not stop taking them two weeks before surgery

5. Patients with previous cervical surgeries, psychiatric disorders and patients receiving opioid analgesic medications within 24 hours preoperatively

#### Date of first enrolment

01/05/2015

Date of final enrolment 01/04/2017

## Locations

**Countries of recruitment** Egypt

#### Study participating centre

Minia University Hospital Corniche El Nil El Corniche El Menia Menia Governorate Minia Egypt 61111

## Sponsor information

## Organisation

Al-Minia University

#### Sponsor details

Faculty of Medicine Main Road Shalaby Land - Menia Al-Minia Egypt 61111 +20 86 23 42584 dean.med@mu.edu.eg

#### Sponsor type

University/education

Website http://www.minia.edu.eg/med/

ROR https://ror.org/02hcv4z63

# Funder(s)

**Funder type** Other

**Funder Name** Investigator initiated and funded

# **Results and Publications**

**Publication and dissemination plan** Publication of the research is planned to be after registering the clinical trial as soon as possible.

Intention to publish date 30/06/2017

#### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from josefzekry2@yahoo.com

#### IPD sharing plan summary

Available on request

Study outputs					
Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	02/12/2017	26/11/2020	Yes	No