Perioperative Duloxetine and Etoricoxib improve postoperative pain

Submission date	Recruitment status No longer recruiting	Prospectively registered	
17/06/2017		☐ Protocol	
Registration date	Overall study status	Statistical analysis plan	
19/06/2017	Completed	[X] Results	
Last Edited	Condition category	Individual participant data	
26/11/2020	Surgery		

Plain English summary of protocol

Background and study aims

Having spinal surgery requires a lot of recovery time. Usually patients who undergo surgery need medication to help them recover and control any after surgery pain. There are a variety of medications that can be used to help treat patients. Some suggestions for medications to use include duloxetine (which is used to treat anxiety, depression, neurological pain) as well as etoricoxib (an anti-inflammatory (swelling) painkiller). It can be sometimes difficult for healthcare workers to figure out what the best combination of medications is most effective and provides patients with the least amount of side effects while still improving their pain symptoms. The aim of this study is to find the best regimen of drugs for the postoperative (after surgery) analgesia (pain prevention) for patients undergoing spinal surgeries in order to improve patient satisfaction of their postoperative care and to provide them with less side effects from medication after the surgery.

Who can participate?

Adults patients aged 18 and older who are having spinal surgery.

What does the study involve?

Participants are randomly allocated to one of four groups. Those in the first group receive a capsule containing a placebo (dummy drug) and two placebo tablets an hour before surgery and then again 24 hours later. Those in the second group receive a capsule a placebo and two etoricoxib tablets an hour before surgery and then again 24 hours later. Those in the first group receive a capsule containing duloxetine and two placebo tablets an hour before surgery and then again 24 hours later. Those in the fourth group receive a capsule containing duloxetine and two etoricoxib tablets an hour before surgery and then again 24 hours later. Patients then have their surgery as per standard practice. Half an hour after their anaesthesia has worn off and then two, four, six, 12, 24 and 48 hours later, participants are asked to rate their pain levels. Their satisfaction with the treatment they received and pain killer use is also recorded.

What are the possible benefits and risks of participating?

Participants may benefit from being more satisfied with their surgery and less side effects from drugs after the surgery. There are no notable risks with participating however there are risks with any medication and this should be monitored.

Where is the study run from? Minia University Hospital (Egypt)

When is the study starting and how long is it expected to run for? April 2015 to May 2017

Who is funding the study? Investigator initiated and funded (Egypt)

Who is the main contact? Mr Josef Zekry Attia

Contact information

Type(s)

Scientific

Contact name

Mr Josef Zekry Attia

ORCID ID

https://orcid.org/0000-0003-4842-231X

Contact details

Faculty of Medicine Al-Minia University Main Road Shalaby Land - Menia Al-Minia Egypt 61111 +20 10 00 427407 josefzekry2@yahoo.com

Additional identifiers

Protocol serial number

7

Study information

Scientific Title

Perioperative Duloxetine and Etoricoxib to improve postoperative pain after lumbar laminectomy: A randomized, double-blind, controlled study

Study objectives

Perioperative Duloxetine and Etoricoxib given together are more effective than when given individually at reducing postoperative pain after lumbar laminectomy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Ethics Committee at Faculty of Medicine Minia university, 01/05/2015, ref: 1-5-2015

Study design

Four-arm randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Postoperative pain after laminectomy surguries

Interventions

Patients admitted to the hospital are randomised using computer-generated random numbers with closed-sealed envelopes into one of the four groups of 30 patients.

Group 1: Participants receive a placebo capsule + two placebo tablets one hour before surgery and again 24 hours later

Group 2: Participants receive a placebo capsule + 90 mg etoricoxib tablets one hour before surgery and again 24 hours later

Group 3: Participants receive a 60 mg duloxetine capsule + two placebo tablets one hour before surgery and again 24 hours later

Group 4: Participants receive a 60 mg duloxetine capsule + two 90 mg etoricoxib tablets one hour before surgery and again 24 hours later

Participants in all groups undergo surgery as per standard practice.

Participants in all groups are asked to rate their pain levels 30 minutes after the end of anesthesia and two, four, six, 12, 24 and 48 hours post-operatively. In addition, patient satisfaction and pain killer use is also assessed 24 and 48 hours after surgery.

Intervention Type

Other

Primary outcome(s)

Pain is measured using pain assessment measures and questions 30 minutes after the end of anesthesia and two, four, six, 12, 24 and 48 hours post-operatively

Key secondary outcome(s))

- 1. The time to first rescue analgesic is routinely 24h and 48h postoperatively
- 2. Total morphine consumption is measured by calculating the doses at 24 and 48 hours postoperatively
- 3. Side effects, including headache, rash, nausea, vomiting, dizziness and drowsiness are recorded. The severity of postoperative nausea and vomiting (PONV) is graded on a four-point ordinal scale

4. Patient satisfaction is measured using a numerical rating scale (NRS) at 24 hours postoperatively

Completion date

01/05/2017

Eligibility

Key inclusion criteria

- 1. Patients with an ASA physical status of I, II and III scheduled for single level lumbar spinal disc prolapsed surgery
- 2. Can provide written informed consent
- 3. Aged between 18 and older

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

ΔII

Total final enrolment

120

Key exclusion criteria

- 1. History of allergic reaction to any of the study drugs
- 2. History of drug or alcohol abuse
- 3. Abnormal renal or liver function tests.
- 4. Patients using antidepressants who did not stop taking them two weeks before surgery
- 5. Patients with previous cervical surgeries, psychiatric disorders and patients receiving opioid analgesic medications within 24 hours preoperatively

Date of first enrolment

01/05/2015

Date of final enrolment

01/04/2017

Locations

Countries of recruitment

Egypt

Study participating centre Minia University Hospital

Corniche El Nil El Corniche El Menia Menia Governorate Minia Egypt 61111

Sponsor information

Organisation

Al-Minia University

ROR

https://ror.org/02hcv4z63

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from josefzekry2@yahoo.com

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added Peer review	ed? Patient-facing?
Results article	results	02/12/2017	26/11/2020 Yes	No

Yes

Participant information sheet 11/11/2025 No