Comparative clinical trial of Krytantek Ofteno® versus Combigan D® in patients with primary open-angle glaucoma

Submission date	Recruitment status	Prospectively registered
04/04/2008	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
17/07/2008	Completed	Results
Last Edited	Condition category	[] Individual participant data
17/07/2008	Eye Diseases	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Leopoldo Baiza-Duran

Contact details

Hidalgo No. 861-A Colonia Centro Sector Hidalgo Guadalajara Mexico 44100 +52 (01)33 3826 4152 drbvista@sophia.com.mx

Additional identifiers

Protocol serial number

KKVSCD0208FII

Study information

Scientific Title

Comparative clinical trial of the safety and efficacy of Krytantek Ofteno® formulated by Laboratorios Sophia S.A. de C.V. versus Combigan D® in the treatment of patients with primary open-angle glaucoma or ocular hypertension

Acronym

KOEG

Study objectives

There is no difference between the hypotensive effect of Krytantek Ofteno® versus Combigan D® in patients with a diagnosis of primary open-angle glaucoma (POAG) and/or ocular hypertension with or without pseudoexfoliation.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the local ethics committee (Nuestra Señora de la Luz Hospital, Mexico City) on the 20th June 2008.

Study design

Multicentre, prospective, longitudinal, comparative, double-blind clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Primary open angle glaucoma

Interventions

According to the random chart, 44 patients will receive over each eye one drop of a topical ophthalmic solution composed of timolol 0.5%, dorzolamide 2%, and brimonidine 0.2% in a fixed-combination formula (Krytantek Ofteno®) developed by Laboratorios Sophia S.A. de C.V. each 12 hours. The other 44 patients will be administered one drop of Combigan D® over each eye every 12 hours; both medications will be administered during a period of 90 days. All the study articles will be labeled with a non-consecutive "code number" that is randomly-generated by computer.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Krytantek Ofteno®, Combigan D®

Primary outcome(s)

Intraocular pressure, measured at days 0, 7, 15, 30, 60 and 90. In all days the measurement will be at 8:00 am.

Key secondary outcome(s))

- 1. Visual fields, evaluation at at days 0 and 90
- 2. Ocular surface fluorescein staining, measured at at days 0 and 90

Completion date

30/04/2009

Eligibility

Key inclusion criteria

- 1. Patients must have a clinical diagnosis of mild to moderate primary open-angle glaucoma (POAG) with or without pseudoexfoliation and pigmentary dispersion or ocular hypertension
- 2. Patients may be of both masculine or feminine genders and must be 18 years old or older
- 3. Patients with intraocular pressure between 21 30 mmHg

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Patients with one blind eye
- 2. Patients with visual acuity of 20/40 or worst in any of both eyes without a justifying cause
- 3. Patients with active stage history of any concomitant ocular disease besides the required one
- 4. Patients taking any medication, whether topically or by any route of administration, that decisively interferes in the study's results, until 48 hours previous to the day 1 of the trial or until a time period in which residual effects could be present
- 5. Sulfa allergy patients
- 6. Patients with history of hypersensibility or any medical situation that contraindicates or makes the using of risky for any of the study articles or their compounds under any route of administration as well as any drug or formulation derived from them or related to them
- 7. Contact lenses users
- 8. Fertile-age women who are not using an adequate contraceptive method as well as pregnant or breast-feeding women
- 9. Patients with history of cataract surgery with or without IOL implant (Pseudoaphakia or aphakia) three months or less prior to day 1 of the trial

- 10. Patients enrolled in any medical trial out of the Laboratorios Sophia S.A. de C.V. sponsorship under the last 90 days prior to this trial
- 11. Legally disqualified or mentally disabled patients who cannot sign the informed consent to participate in this clinical trial
- 12. Patients who cannot comply with the medical appointments or with all the protocol requirements
- 13. Patients who refuse to participate in this clinical trial
- 14. Patients with optic disc excavation equal to 0.8 or more
- 15. Normal-tension glaucoma patients

Date of first enrolment 30/04/2008

Date of final enrolment 30/04/2009

Locations

Countries of recruitment Bolivia
Chile
Colombia
Dominican Republic
Ecuador
El Salvador
Guatemala
Honduras
Mexico
Nicaragua
Panama
Peru

Study participating centre Hidalgo No. 861-A Guadalajara Mexico 44100

Venezuela

Sponsor information

Organisation

Laboratorios Sophia S.A. de C.V. (Mexico)

ROR

https://ror.org/00zpf2822

Funder(s)

Funder type

Industry

Funder Name

Laboratorios Sophia S.A. de C.V. (Mexico)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes