High concentration oxygen therapy in pediatric asthma exacerbation

Submission date	Recruitment status	Prospectively registered
15/06/2016	No longer recruiting	Protocol
Registration date 17/06/2016	Overall study status Completed	Statistical analysis plan
		Results
Last Edited	Condition category	Individual participant data
17/05/2017	Respiratory	Record updated in last year

Plain English summary of protocol

Background and study aims

Asthma Is a common long term health condition caused by inflammation of the bronchi, small tubes that carry air in and out of the lungs. The airways can become narrow if the sufferer comes into something that "triggers" their asthma – for example animal fur, pollen, viral infection and exercise. This can cause symptoms such as coughing, wheezing, tightness of the chest and feeling breathless. Occasionally, asthma symptoms can gradually or suddenly get much worse. This is called an asthma attack. Severe attacks may require hospital treatment and, in extreme cases, can even lead to death. This study is looking at the use of oxygen in children who are having an asthma attack. The researchers are investigating whether the amount of oxygen used affects the carbon dioxide in children suffering from an asthma attack and whether it makes their asthma worse and/or results in their hospitalization.

Who can participate?

Children 2-18 years of age suffering from a moderate to severe asthma attack and are coming to the emergency room (ED)

What does the study involve?

Participants are randomly allocated to one of two groups. Those in group 1 receive 100% oxygen as part of their treatment. Those in group 2 receive titrated oxygen to keep that oxygen level between 92 to 95% while they are receiving their nebulization treatments.

What are the possible benefits and risks of participating?

There are no direct benefits from participating in this study. There are no risks from participating in this study as the oxygen levels are always kept in the normal range in both groups.

Where is the study run from?

The Children's Hospital at Montefiore, Bronx, New York (USA)

When is study starting and how long is it expected to run for? December 2013 to May 2016

Who is funding the study?
Pediatric Critical Care Division, Children's Hospital at Montefiore (USA)

Who is the main contact? Dr Bhavi Patel

Contact information

Type(s)

Public

Contact name

Dr Bhavi Patel

Contact details

The Children's Hospital at Montefiore 3415 Bainbridge Avenue Bronx New York United States of America 10467

Additional identifiers

Protocol serial number

2015-4431

Study information

Scientific Title

Randomized clinical trial of high concentration oxygen therapy versus titrated oxygen therapy in pediatric asthma exacerbation

Study objectives

High concentration oxygen leads to rise in transcutaneous carbon dioxide in acute exacerbation of asthma in the pediatric population.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Albert Einstein College of Medicine Institutional Board Review, 22/04/2015, ref: 2015-4431

Study design

Single-center randomized prospective double-blind clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pediatric asthma exacerbation

Interventions

Children 2 to 18 years with previously diagnosed asthma with moderate to severe asthma exacerbation (asthma score > 5) were randomized to high concentration oxygen therapy (100% oxygen via face mask at >4L/min) or titrated oxygen therapy (titrated up from 21% via a blender continuously) to maintain saturations between 92 to 95% while receiving their nebulized treatments.

Randomization was based on computer randomization sequence program generated by the biostatistician.

Intervention Type

Other

Primary outcome(s)

Rise in transcutaneous carbon dioxide, measured via a transcutaneous carbon dioxide monitor at 0, 20, 40, 60 minutes and then every 30 minutes till disposition decision was reached by the ED physician

Key secondary outcome(s))

- 1. Change of > 4 mmHg PtCO2 and > 38 mmHg PtCO2 measured via a transcutaneous Co2 monitor at 0, 20, 40, 60 minutes and then every 30 minutes till disposition decision was reached by the ED physician
- 2. Change in asthma score, measured via a validated respiratory score tool used by the hospital as part of the asthma treatment pathway at 0, 20, 40, 60 minutes and then every 30 minutes till disposition decision was reached by the ED physician
- 3. Admission rates

Completion date

18/05/2016

Eligibility

Key inclusion criteria

Patients between 2 to 18 years with previous medical diagnosis of asthma presenting with moderate to severe asthma exacerbation with asthma score > 5

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

2 years

Upper age limit

18 years

Sex

All

Key exclusion criteria

- 1. Disorders with hypercapnic respiratory failure such as neuromuscular disease, chest wall restriction, or obesity hypoventilation
- 2. Unconscious patient
- 3. Pregnancy
- 4. Receiving sedatives/depressants
- 5. History of congenital heart disease
- 6. Disease of cardiac, renal, or hepatic systems
- 7. History of smoking

Date of first enrolment

01/08/2015

Date of final enrolment

18/05/2016

Locations

Countries of recruitment

United States of America

Study participating centre The Children's Hospital at Montefiore

3415 Bainbridge Avenue Bronx New York United States of America 10467

Sponsor information

Organisation

Pediatric Critical Care Department

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Department of Pediatric Critical Care at the Children's Hospital at Montefiore (CHAM)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing? Participant information sheet 11/11/2025 No

Yes