# Comparison of different techniques regarding visualization of scars in the heart muscle using cardiac MRI

Submission date	Recruitment status	Prospec
06/06/2017	No longer recruiting	[] Protoco
Registration date	Overall study status	[] Statistic
10/07/2017	Completed	[X] Results
Last Edited 15/03/2019	<b>Condition category</b> Circulatory System	[_] Individu

- ctively registered
- ol
- ical analysis plan
- ual participant data

#### Plain English summary of protocol

#### Background and study aims

A cardiac MRI scan is a non-invasive test where magnetic and radio waves are used to create pictures of the inside of the heart. Over the last couple of years cardiac MRI has become the gold standard method for looking at scars in the heart muscle. This scarring can be due to myocardial infarction (heart attack), thickening of the heart walls or inflammation of the heart muscle itself and is a very strong indicator of the likely course of these conditions. Cardiac MRI uses a technique called late gadolinium enhancement (LGE) to show scars. The gold standard LGE technique is a so-called segmented PSIR sequence and provides great image guality, but only in patients with a stable heart rhythm and able to hold their breath, which is required for the technique to work. Unfortunately, many patients do not meet these requirements. New socalled multi-slice LGE sequences are faster techniques than the gold standard method and also work for patients with irregular heart rhythm or shortness of breath. The aim of this study is to assess the accuracy of three of these novel multi-slice LGE sequences to detect and quantify scars in the heart muscle when compared to a gold standard PSIR sequence.

#### Who can participate?

Patients aged 18 and over referred for cardiac MRI after a heart attack (myocardial infarction), with thickened heart walls (hypertrophic cardiomyopathies), or due to suspected or known inflammation of the heart muscle cells

#### What does the study involve?

All participants undergo a standard cardiac MRI scan with the gold standard LGE PSIR sequence and also the new multi-slice sequences. This prolongs the MRI scan by about five minutes. The size of the myocardium (heart muscle) and image quality are compared between the scans.

#### What are the possible benefits and risks of participating?

Possible benefits include improved detection of scars in the heart muscle. If the multi-slice sequences are found to perform as well as the gold standard sequence, these faster sequences could be used in future to improve access to cardiac MRI by decreasing scan time at a high image quality and increasing patient comfort. As all patients receive the cardiac MRI even if they do not participate and the scan itself is only prolonged by 5 minutes (of a 45 minutes total scan time), there is no additional risk or burden expected with participation in this study.

Where is the study run from? HELIOS Clinic Berlin-Buch (Germany)

When is the study starting and how long is it expected to run for? November 2014 to December 2016

Who is funding the study? Charité Universitätsmedizin Berlin (Germany)

Who is the main contact? Prof. Jeanette Schulz-Menger

# **Contact information**

**Type(s)** Scientific

**Contact name** Prof Jeanette Schulz-Menger

#### **Contact details**

Charité University Medicine Berlin Campus Buch Working Group Kardiale MRT Lindenberger Weg 80 Berlin Germany 13125

## Additional identifiers

EudraCT/CTIS number

**IRAS number** 

ClinicalTrials.gov number

**Secondary identifying numbers** LGE COMPARE (internal study code)

# Study information

#### Scientific Title

Comparison of fast multi-slice and standard segmented techniques for detection of late gadolinium enhancement in ischemic and non-ischemic cardiomyopathy - a prospective clinical trial

#### **Study objectives**

All tested fast multi-slice LGE sequences are non-inferior to the segmented PSIR gold standard method with regard to LGE size quantification (in gram) using a semi-automated threshold method.

### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Ethics board at Charité University Medicine Berlin, Campus Mitte, 16/10/2014, ref: EA1/305/14

**Study design** Observational cohort study

**Primary study design** Observational

**Secondary study design** Cohort study

**Study setting(s)** Hospital

**Study type(s)** Diagnostic

#### Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

#### Health condition(s) or problem(s) studied

Myocardial infarction, hypertrophic cardiomyopathy and inflammatory heart disease

#### Interventions

This is an observational study in which three different techniques are compared for visualization of myocardial fibrosis using late Gadolinium enhancement in patients with known or suspected cardiomyopathy.

Patients sent for clinically indicated cardiac LGE-based MRI with known or suspected LGE due to chronic infarction, inflammatory myocardial disease and hypertrophic cardiomyopathy (HCM) are prospectively recruited. The indication for the MRI itself is clinical. However, for this study the MRI protocol is extended by the tested multislice LGE sequences for approximately 5 extra minutes. LGE images are acquired using three different LGE sequences. All patients were scanned with all three LGE sequences (intraindividual comparison). All LGE sequences are then assessed as follows: image quality is evaluated with a 4-point scoring system. Contrast-to-noise ratios and acquisition time are measured. Size of LGE is quantitatively assessed using a semi-automated threshold method. All assessments are done by blinded readers.

#### Intervention Type

Other

#### Primary outcome measure

Size of myocardium with late Gadolinium enhancement in each tested MR sequence, measured by semiautomated threshold method in MRI DICOM data at one timepoint, the date of the MRI (no follow up)

#### Secondary outcome measures

1. Contrast-to-noise ratio in each tested MR sequence, measured using MRI DICOM data 2. Image quality, scored on a 4-point-scaling system based on visual assessment by blinded readers

Measured at at one timepoint, the date of the MRI (no follow up)

#### Overall study start date

01/11/2014

#### **Completion date**

31/12/2016

# Eligibility

#### Key inclusion criteria

1. Known or suspected chronic myocardial infarction, hypertrophic cardiomyopathy or inflammatory heart disease

- 2. Clinical indication for MR exam with late gadolinium enhancement
- 3. Age range ≥18 years (no upper limit)

### Participant type(s)

Patient

### Age group

Adult

### Lower age limit

18 Years

#### Sex

Both

# **Target number of participants** 302

#### Key exclusion criteria

Any contraindication for MR exam
Acute or chronic renal failure with GFR <30ml/min</li>

### Date of first enrolment

01/11/2014

### Date of final enrolment

31/10/2016

## Locations

**Countries of recruitment** Germany

**Study participating centre HELIOS Clinic Berlin-Buch** Germany 13125

# Sponsor information

**Organisation** Charité University Medicine Berlin

**Sponsor details** Working Group Kardiale MRT Lindenberger Weg 80 Berlin Germany 13125

**Sponsor type** University/education

Website http://www.cmr-berlin.org

ROR https://ror.org/001w7jn25

# Funder(s)

**Funder type** University/education

**Funder Name** Charité Universitätsmedizin Berlin

Alternative Name(s) Medical School - Charité - University Medicine Berlin **Funding Body Type** Private sector organisation

#### Funding Body Subtype

For-profit companies (industry)

### Location

Germany

# **Results and Publications**

#### Publication and dissemination plan

Results of this study shall be published in a high-ranking peer-reviewed journal

#### Intention to publish date

31/12/2017

#### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study is not expected to be made available due to reasons of data protection laws in Germany. However, upon request methodology and data set structure can be shared.

#### IPD sharing plan summary

Not expected to be made available

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	19/02/2018	15/03/2019	Yes	No