Hysterectomy or Endometrial AbLation Trial for Heavy menstrual bleeding

Submission date	Recruitment status No longer recruiting	Prospectively registered	
27/01/2014		☐ Protocol	
Registration date	Overall study status	Statistical analysis plan	
28/01/2014	Completed	[X] Results	
Last Edited 19/05/2023	Condition category Urological and Genital Diseases	[] Individual participant data	

Plain English summary of protocol

Background and study aims

Heavy menstrual bleeding (HMB) is a very common and distressing condition which affects over 1.5 million women in the UK. It is responsible for one in 20 women consulting their general practitioner (GP) and accounts for 20% of hospital referrals in gynaecology. Initial treatment usually involves the use of drugs, including the combined oral contraceptive pill, as well as a hormone-impregnated intrauterine contraceptive device. However, these may be unsuitable or unsuccessful in some women, who will need surgical treatment. The main aim of this study is to compare the clinical and cost effectiveness of two different surgical treatments: laparoscopic supra-cervical hysterectomy (LASH) and endometrial ablation (EA) for the treatment of HMB.

Who can participate?

Women less than 50 years of age with heavy menstrual bleeding (HMB) eligible for surgical treatment can take part.

What does the study involve?

Eligible women will be randomly allocated to undergo either LASH or EA. All participants will be asked to complete assessment questionnaires at certain time points - before surgery and again at 6 weeks, 6 and 12 months after surgery. Women will also complete a 14-day diary immediately after their operation and a simple questionnaire at 4 weeks after surgery. Medium-term follow-up involves the completion of an additional questionnaire at 5-8 years post-randomisation (as of 15/05/2023).

What are the possible benefits and risks of participating?

Participants may not benefit personally from taking part in the study but will be directly helping us to generate information which could help plan more effective treatment for women with heavy periods. We do not think that there are any additional risks or disadvantages to participating in this study. Whichever group participants are allocated to, their care will be overseen by an experienced consultant gynaecologist. Steps are always taken to make sure that any possible risks are minimised. As part of routine care, participants will be well informed of potential risks.

Where is the study run from?

Patients will be recruited from approximately 26 NHS hospitals in the UK:

Aberdeen Royal Infirmary, Aberdeen; Forth Valley Royal Hospital, Larbert; Countess of Chester Hospital NHS Foundation Trust, Chester; Castle Hill Hospital, Cottingham; Queen Elizabeth Hospital, Edgbaston, Birmingham; Gartnavel Royal Hospital, Glasgow; Sunderland Royal Hospital, Sunderland; The Royal Victoria Infirmary, Newcastle Upon Tyne; Harrogate District Hospital, Harrogate; Northern General Hospital, Sheffield; Worcester Royal Hospital, Worcester; Arrowe Park Hospital, Upton, Wirral; Stepping Hill Hospital, Stockport; Singleton Hospital, Swansea; Royal Cornwall Hospital, Cornwall; Queen Alexandra Hospital, Portsmouth; The Royal Hampshire County Hospital, Winchester; Whipps Cross Hospital, London; Princess Royal University Hospital, Farnborough Common, Kent; Derriford Hospital, Plymouth; The Great Western Hospital, Swindon; St Peter's Hospital, Chertsey, Surrey; Princess Anne Hospital, Southampton; Royal Sussex County Hospital, Brighton; Poole Hospital NHS Foundation Trust; Poole; Worthing Hospital, Worthing.

When is the study starting and how long is it expected to run for? January 2014 to August 2024 (as of 15/05/2023)

Who is funding the study? National Institute for Health Research (NIHR), UK.

Who is the main contact? Centre for Healthcare Randomised Trials chart@abdn.ac.uk

Contact information

Type(s)

Scientific

Contact name

Prof Siladitya Bhattacharya

Contact details

School of Medicine and Dentistry University of Aberdeen Aberdeen Maternity Hospital 2nd Floor, Foresterhill Aberdeen United Kingdom AB25 2ZD

s.bhattacharya@abdn.ac.uk

Additional identifiers

Protocol serial number 3/073/13

Study information

Scientific Title

Hysterectomy or Endometrial AbLation Trial for Heavy menstrual bleeding. A multicentre randomised controlled trial comparing laparoscopic supra-cervical hysterectomy with second generation endometrial ablation for the treatment of heavy menstrual bleeding

Acronym

HEALTH

Study objectives

The hypothesis being tested is that laparoscopic supra-cervical hysterectomy is superior to second generation endometrial ablation for the treatment of heavy menstrual bleeding (HMB) in terms of patient satisfaction, quality of life (QoL) and costs.

Ethics approval required

Old ethics approval format

Ethics approval(s)

North of Scotland Research Ethics Service Committee 2, 06/01/2014, ref. 13/NS/0155

Study design

Multicentre randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Heavy menstrual bleeding

Interventions

The patients are randomised to two groups.

The interventions being compared are:

1. Laparoscopic supra-cervical hysterectomy (LASH):

The LASH procedure involves removing the part of your womb that causes menstrual bleeding. This is done by keyhole surgery. Three small cuts on the stomach wall are made. The procedure is done under a general anaesthetic and patients are usually home within 24 hours of the operation. As the cervix is not removed, women still need to have cervical smears in the future.

2. Endometrial ablation (EA): The EA procedure involves placing a thin device in the womb by passing it first through the vagina and then through the cervix. The lining of the womb is destroyed and the device is then removed. EA is done as a day case procedure in hospital. It is usually done under a general anaesthetic but can be under local anaesthetic if preferred. As the uterus is not removed women still need cervical smears in the future.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

- 1. Menorrhagia multi-attribute scale (MMAS), a condition-specific Quality of Life outcome
- 2. Patient satisfaction, measured on a six point scale (from totally satisfied to totally dissatisfied) measured at 12 months post surgery
- 3. Incremental cost (to the health service) per quality-adjusted life year (QALY) gained (LASH versus EA)

Key secondary outcome(s))

Current secondary outcome measures as of 15/05/2023:

Patient-reported:

- 1. MMAS at 6 months and 5-8 years post-randomisation
- 2. Patient-reported satisfaction at 6 months and 5-8 years post-randomisation
- 3. Acceptability of procedure measured at 6 weeks
- 4. Severity of postoperative pain using a pain Numerical Rating Scale (NRS) measured at 1-14 days and at 6 weeks, symptom diary days 1 to 14 (including analgesic use)
- 5. Generic health-related quality of life (SF-12, EQ-5D 3-L) measured at baseline, 6 months, 12 months and 5-8 years post-randomisation
- 6. Sexual Activity Questionnaire (SAQ) at baseline, 6 and 12 months

Clinical

- 1. Duration of operation
- 2. Peri-operative complications and recovery details including analgesia requirements
- 3. Time to discharge
- 4. Further gynaecological surgery in 12 months and 5-8 years post-randomisation

Economic

Wider societal costs associated with changes in productivity based on information on the time taken to return to normal activities (following intervention) combined with questions on work productivity delivered during the follow-up period. Further, a simple Markov model, based on within trial data supplemented by available published data on the requirement for further gynaecological surgery over time (following the alternative procedures) will be developed and used to extrapolate cost-effectiveness beyond 12 months.

Previous secondary outcome measures:

Patient reported:

- 1. MMAS at 6 months
- 2. Patient reported satisfaction at 6 months
- 3. Acceptability of procedure measured at 6 weeks
- 4. Severity of post operative pain using a pain Numerical Rating Scale (NRS) measured at 1-14 days and at 6 weeks, symptom diary days 1 to 14 (including analgesic use)
- 5. Generic health related quality of life (SF-12, EQ-5D 3-L) measured at baseline, 6 and 12 months
- 6. Sexual Activity Questionnaire (SAQ) at baseline, 6 and 12 months

Clinical

- 1. Duration of operation
- 2. Peri-operative complications and recovery details including analgesia requirements
- 3. Time to discharge
- 4. Further gynaecological surgery by 12 months

Economic

Wider societal costs associated with changes in productivity based on information on the time

taken to return to normal activities (following intervention) combined with questions on work productivity delivered during the follow-up period. Further, a simple Markov model, based on within trial data supplemented by available published data on the requirement for further gynaecological surgery over time (following the alternative procedures) will be developed and used to extrapolate cost-effectiveness beyond 12 months.

Completion date

31/08/2024

Eligibility

Key inclusion criteria

- 1. Women less than 50 years of age with heavy menstrual bleeding eligible for endometrial ablation
- 2. Women who are willing to be randomised between laparoscopic supra-cervical hysterectomy and endometrial ablation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

660

Key exclusion criteria

- 1. Women with plans to conceive, endometrial atypia, uterine cavity size greater than 11 cm, submucosal fibroids distorting the uterine cavity, contradictions for laparoscopic surgery (e.g. midline lower abdominal incision or known intrabdominal / pelvic adhesions) and previous endometrial ablation (EA)
- 2. Women who are unable to give informed consent or complete trial documentation

Date of first enrolment

01/01/2014

Date of final enrolment

31/03/2017

Locations

Countries of recruitment

United Kingdom

Study participating centre School of Medicine and Dentistry Aberdeen United Kingdom AB25 2ZD

Sponsor information

Organisation

University of Aberdeen/NHS Grampian (UK)

ROR

https://ror.org/016476m91

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research (NIHR) (UK) - NIHR Evaluation, Trials and Studies Coordinating Centre (NETSCC) - NIHR Health Technology Assessment Programme (HTA), ref: 12/35/23

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from the Chief Investigator, Prof Siladitya Bhattacharya, s.bhattacharya@abdn.ac. uk.

IPD sharing plan summary

Available on request

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Results article results 19/10/2019 17/09/2019 Yes No

results

Results article		01/09/2019	03/10/2019 Yes	No
HRA research summary			28/06/2023 No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025 No	Yes
Study website	Study website	11/11/2025	11/11/2025 No	Yes