# Comparison of minimally invasive versus conventional laparoscopic (keyhole) bariatric surgery

Submission date	Recruitment status	[X] Prospectively registered
19/02/2021	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
04/03/2021	Completed	Results
Last Edited	Condition category	Individual participant data
05/07/2024	Surgery	<ul><li>Record updated in last year</li></ul>

# Plain English summary of protocol

Background and study aims

Laparoscopic Sleeve Gastrectomy (also called bariatric surgery) is the most performed bariatric surgery worldwide with 155,000 procedures in the United States in 2018 and 36,000 in France in 2016.

LSG surgery is done in a hospital under general anesthesia. The surgeon will make about five small cuts in the belly. He or she will do the surgery using a thin, long, telescope with a tiny camera at the end. Instruments pushed through the incisions will be used to remove about 80% of the stomach. The surgeon will do the procedure using images on a TV screen in the operating room.

This surgery takes out the part of the stomach that curves outward, called the fundus. After the fundus is taken out, the surgeon will close the rest of the stomach into a tube shape that looks like a banana or the sleeve of the shirt, hence the name "sleeve gastrectomy."

A recent systematic review demonstrated the safety and efficacity of single incision laparoscopic sleeve gastrectomy. But to our knowledge, no trial has compared traditional multiport sleeve gastrectomy to single incision laparoscopic sleeve gastrectomy. The aim of this study is to compare these two techniques and to determine the superiority of one of them considering the following aspects: postoperative pain, postoperative scars esthetic aspect, postoperative sex life along with postoperative diurnal and nocturnal quality of life.

## Who can participate?

Adult patients with body mass index >40 kg/m<sup>2</sup> or with a body mass index >35 kg/m<sup>2</sup> with serious comorbidity such as obstructive sleep apnea, diabetes mellitus or hypertension undergoing laparoscopic sleeve gastrectomy as a primary bariatric procedure.

#### What does the study involve?

As participants undergoing sleeve gastrectomy (regardless of number of used ports) sign up for a scheduled postoperative follow-up, no additional outpatient visit or examination is required in this study.

What are the possible benefits and risks of participating?

This study is a non-randomized trial. Patients will be freely addressed to one of two teams of surgeons who adopted and routinely perform exclusively one of two surgical approaches (conventional laparoscopy or single-port laparoscopic sleeve gastrectomy). Therefore, no specific additional benefits and risks are expected from the participation of patients in the trial.

Where is the study run from? Antoine Beclere Hospital of Assistance Publique-Hôpitaux de Paris (France)

When is the study starting and how long is it expected to run for? January 2021 to June 2024

Who is funding the study? Investigator initiated and funded

Who is the main contact? Dr Hadrien Tranchart hadrien.tranchart@aphp.fr

# Contact information

# Type(s)

Public

#### Contact name

Dr Joseph Derienne

#### **ORCID ID**

https://orcid.org/0000-0002-2562-6507

#### Contact details

Department of minimally invasive digestive surgery Hôpital Antoine Béclère 157 rue de la porte de trivaux Clamart France 92140 +33 (0)1 45 37 43 47 joseph.derienne@aphp.fr

# Type(s)

Scientific

#### Contact name

Dr Hadrien Tranchart

#### **ORCID ID**

https://orcid.org/0000-0003-2173-2828

#### Contact details

Department of minimally invasive digestive surgery Hôpital Antoine Béclère 157 rue de la porte de trivaux Clamart France 92140 +33 (0)1 45 37 43 47 hadrien.tranchart@aphp.fr

# Additional identifiers

# Clinical Trials Information System (CTIS)

Nil known

# ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

Single incision laparoscopic sleeve gastrectomy versus conventional multiport sleeve gastrectomy (ONE SLEEVE): a prospective non-randomized controlled trial

#### Acronym

**ONE SLEEVE** 

# **Study objectives**

Single incision laparoscopic sleeve gastrectomy provides an improved postoperative outcome compared to conventional multiport sleeve gastrectomy.

# Ethics approval required

Old ethics approval format

# Ethics approval(s)

No ethics approval is required for this study, since it is a nonrandomized controlled trial of a routine procedure. This was validated with the clinical research unit of the Paris-Sud University.

# Study design

Single-centre prospective non-randomized controlled observational study

# Primary study design

Observational

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Laparoscopic sleeve gastrectomy

#### **Interventions**

Patients undergo a laparoscopic sleeve gastrectomy either using a single incision or a conventional multiport technique depending on the surgical team they are assigned to. This is a routine procedure.

There is a one-year follow-up which is not additional to usual treatment.

# Intervention Type

Procedure/Surgery

# Primary outcome(s)

A written questionnaire with 0-10 scales is completed at 3 months evaluating:

- 1. Scar esthetic impact on patients
- 2. Postoperative parietal pain
- 3. Impact of the scar on postoperative sex life
- 4. Scar impact on diurnal quality of life (daily basis movements, work)
- 5. Scar impact on nocturnal quality of life (sleep)

# Key secondary outcome(s))

- 1. 6 and 12 months written questionnaires with 0-10 scales evaluating:
- 1.1. Scar esthetic impact on patients
- 1.2. Postoperative parietal pain
- 1.3. Impact of the scar on postoperative sex life
- 1.4. Scar impact on diurnal quality of life (daily basis movements, work)
- 1.5. Scar impact on nocturnal quality of life (sleep)
- 2. 90 days postoperative Morbi-mortality from patient records
- 3. Bodyweight (kg) (at baseline, 3, 6 and 12 months)
- 4. Body Mass index  $(kg/m^2)$  (at baseline, 3, 6 and 12 months)
- 5. Co-morbidities (at baseline and 12 months) from patient records
- 6. Incisional hernia rate (at 12 months) with clinical and radiological (CT scan) evaluation

## Completion date

01/06/2024

# **Eligibility**

#### Key inclusion criteria

- 1. Consecutive patients undergoing laparoscopic sleeve gastrectomy
- 2. Decision for intervention after multidisciplinary discussion
- 3. Sleeve gastrectomy as a primary bariatric procedure
- 4. Body mass index >40 kg/m $^2$  or >35 kg/m $^2$  with severe comorbidities.

# Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

#### Sex

All

#### Total final enrolment

176

# Key exclusion criteria

- 1. Previous upper abdominal surgery (except laparoscopic cholecystectomy)
- 2. Patient under guardianship and trusteeship

## Date of first enrolment

01/06/2021

#### Date of final enrolment

01/06/2023

# Locations

## Countries of recruitment

France

# Study participating centre Hôpital Antoine Béclère

Department of minimally invasive digestive surgery 157 rue de la porte de trivaux Clamart France 92140

# Sponsor information

# Organisation

Hôpital Antoine-Béclère

#### **ROR**

https://ror.org/04sb8a726

# Funder(s)

# Funder type

#### **Funder Name**

Investigator initiated and funded

# **Results and Publications**

# Individual participant data (IPD) sharing plan

All data generated or analysed during this study will be included in the subsequent results publication.

# IPD sharing plan summary

Other

## **Study outputs**

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 11/11/2025 No Yes